Asia-Pacific Symposium for Lattice Field Theory (APLAT 2020)

Contribution ID: 4

Type: not specified

## Study of the rho resonance using the HAL QCD method

*Tuesday, 4 August 2020 15:20 (20 minutes)* 

In this talk, we show the recent status of the rho resonance study in the HAL QCD method. We investigate the I=1 two-pion potential at  $m_{\pi} \approx 411$  MeV by using a new calculation strategy, namely the combination of three techniques: the one-end trick, the sequential propagator, and covariant approximation averaging (CAA).

Thanks to the new strategy, we determine the non-local I=1 two-pion potential at the next-leading-order of the derivative expansion for the first time and obtain the pole of the S-matrix corresponding to the rho resonance. As regards the resonance parameters, our resonance mass is consistent with the previous study using the finite-volume method, but slightly larger decay width and coupling are obtained. We discuss possible origins of this difference.

Primary author: AKAHOSHI (\*), Yutaro (Yukawa Institute for Theoretical Physics, Kyoto University)

Co-authors: HAL QCD COLLABORATION; AOKI, Sinya

Presenter: AKAHOSHI (\*), Yutaro (Yukawa Institute for Theoretical Physics, Kyoto University)

Session Classification: Hadron Spectroscopy and Interactions

Track Classification: Hadron Spectroscopy and Interactions