

# Hadronic light-by-light contribution to $(g - 2)_\mu$ near the SU(3)-flavor-symmetric point

Wednesday, August 5, 2020 2:20 PM (20 minutes)

Understanding the tension between the Standard Model prediction and the experimental results on the anomalous magnetic moment of the muon ( $a_\mu$ ) has been an active research field over the past two decades. The theoretical uncertainty mainly comes from the hadronic contributions, among which the hadronic light-by-light scattering ( $a_\mu^{hlbl}$ ) process plays an important role. We investigate this contribution on the lattice with a position-space approach. In our setup, we treat separately the QED part in the continuum and infinite volume and the QCD part on the lattice, which helps to avoid finite-size effects due to the photon in finite-volume. However, noticeable finite-size effects due to long-distance physics still persist in our approach. In our recent work [arXiv:2006.16224], we have performed computations of  $a_\mu^{hlbl}$  on the lattice with  $N_f = 3$  ensembles having degenerate light and strange quark masses. Our results have satisfactory statistical errors and allow us to concentrate on our strategy to control the finite-size effects using the pion-exchange contribution. The extension of our setup to include non flavor-symmetric ensembles does not require much effort, thanks to the pre-computed single-propagator trace in position space shared among other Mainz projects. Our work in progress toward the physical point will be briefly reported, with focus on the Wick-contraction topologies that vanish at the flavor-symmetric point.

**Primary author:** CHAO (\*), En-Hung (Johannes Gutenberg-Universität Mainz)

**Co-authors:** GÉRARDIN, Antoine; GREEN, Jeremy; HUDSPITH, Renwick J.; MEYER, Harvey B.; OTTNAD, Konstantin

**Presenter:** CHAO (\*), En-Hung (Johannes Gutenberg-Universität Mainz)

**Session Classification:** Hadron Structure

**Track Classification:** Hadron Structure