

Parton distribution functions of Δ^+ on the lattice

Friday, August 7, 2020 5:40 PM (20 minutes)

We perform a first calculation for the unpolarized parton distribution function of the Δ^+ baryon using lattice QCD simulations within the framework of Large Momentum Effective Theory. Two ensembles of $N_f = 2+1+1$ twisted mass fermions are utilized with a pion mass of 270 MeV and 360 MeV, respectively. The baryon, which is treated as a stable single-particle state, is boosted with momentum P_3 with values $\{0.42, 0.83, 1.25\}$ GeV, and we utilize momentum smearing to improve the signal. The unpolarized parton distribution function of Δ^+ is obtained using a non-perturbative renormalization and a one-loop formula for the matching, with encouraging precision. In particular, we compute the $\bar{d}(x) - \bar{u}(x)$ asymmetry and compare it with the same quantity in the nucleon, in a first attempt towards resolving the physical mechanism responsible for generating such asymmetry.

Primary author: LI (*), Yuan (Peking University)

Co-authors: CHAI, Yahui; XIA, Shicheng; ALEXANDROU, Constantia; CICHY, Krzysztof; CONSTANTINOU, Martha; FENG, Xu; HADJIYIANNAKOU, Kyriakos; JANSEN, Karl; KOUTSOU, Giannis; LIU, Chuan; SCAPELLATO, Aurora; STEFFENS, Fernanda

Presenter: LI (*), Yuan (Peking University)

Session Classification: Hadron Structure

Track Classification: Hadron Structure