

Complex Langevin analysis of four-dimensional SU(2) gauge theory with a theta term

Wednesday, August 5, 2020 2:40 PM (20 minutes)

The Monte Carlo simulation of the gauge theory with a theta term is difficult due to the sign problem. We use the complex Langevin method to overcome the problem. In our previous work on two-dimensional U(1) gauge theory with a theta term, we were able to reproduce the exact solution by introducing a puncture on the torus. We also proved that the effect of the puncture disappears in the infinite volume limit as long as $|\theta| < \pi$. In this study, we extend this method to four-dimensional SU(2) gauge theory. Recently the analytic study of 't Hooft anomaly matching condition predicted two possible phase structures around $\theta = \pi$ for this theory. We discuss the possibility of investigating the phase structure by the complex Langevin simulation.

Primary author: MATSUMOTO (*), Akira (SOKENDAI, KEK)

Co-authors: HATAKEYAMA, Kohta; HIRASAWA, Mitsuki; HONDA, Masazumi; ITO, Yuta; NISHIMURA, Jun; YOSPRAKOB, Atis

Presenter: MATSUMOTO (*), Akira (SOKENDAI, KEK)

Session Classification: QCD at nonzero Temperature and Density

Track Classification: QCD at nonzero Temperature and Density