

Long-distance contribution to neutrinoless double beta decays in pion sector

Wednesday, 5 August 2020 14:20 (20 minutes)

Neutrinoless double beta decay, if detected, would prove that neutrinos are Majorana fermions and provide the direct evidence for lepton number violation. If such decay would exist in nature, then $\pi^-\pi^- \rightarrow ee$ and $\pi^- \rightarrow \pi^+ee$ (or equivalently $\pi^-e^+ \rightarrow \pi^+e^-$) are the two simplest processes accessible via first-principle lattice QCD calculations. In this work, we calculate the long-distance contributions to the $\pi^- \rightarrow \pi^+ee$ transition amplitude using four ensembles at the physical pion mass with various volumes and lattice spacings. We adopt the infinite-volume reconstruction method to control the finite-volume effects arising from the (almost) massless neutrino. Providing the lattice QCD inputs for chiral perturbation theory, we obtain the low energy constant $g\pi\pi\nu(m\pi) = -10.89(28)\text{stat}(74)\text{sys}$, which is close to $g\pi\pi\nu(m\pi) = -11.96(31)\text{stat}$ determined from the crossed channel $\pi^-\pi^- \rightarrow ee$ decay.

Primary author: TUO (*), Xin-yu (Peking university)

Co-authors: FENG, Xu; JIN, Lu-chang

Presenter: TUO (*), Xin-yu (Peking university)

Session Classification: Weak Decays and Matrix Elements

Track Classification: Weak Decays and Matrix Elements