


KAON 2022, September 16th 2022

$K \rightarrow \mu^+ \mu^-$
as a third kaon

golden mode

Avital Dery

AD, Ghosh, Grossman, Schacht,
<https://arxiv.org/pdf/2104.06427.pdf>



The message of this talk:

A measurement of time dependence in $K \rightarrow \mu^+ \mu^-$ is
sensitive to the CKM parameter $\bar{\eta}$ with theory
uncertainty of $\mathcal{O}(1\%)$

$$K_L \rightarrow \pi^0 \nu \bar{\nu}$$

$$K^+ \rightarrow \pi^+ \nu \bar{\nu}$$

Theoretically **clean**, sensitive to functions of $V_{ts}^* V_{td}$, $V_{cs}^* V_{cd}$.


What about $K_{L,S} \rightarrow \mu^+ \mu^-$?

- Experimentally straightforward
- Similar weak Hamiltonian (also sensitive to $V_{ts}^* V_{td}$, $V_{cs}^* V_{cd}$)

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \frac{\alpha}{2\pi \sin^2 \Theta_W} \sum_{l=e,\mu,\tau} \left(V_{cs}^* V_{cd} X_{NL}^l + V_{ts}^* V_{td} X(x_t) \right) (\bar{s}d)_{V-A} (\bar{\nu}_l \nu_l)_{V-A}$$


$$K_L \rightarrow \pi^0 \nu \bar{\nu}$$

- Within the SM determined entirely from the weak effective Hamiltonian (*short-distance physics*)
- Purely **CP-violating**




$$K_L \rightarrow \pi^0 \nu \bar{\nu}$$

- Within the SM determined entirely from the weak effective Hamiltonian (**short-distance physics**)
- Purely **CP-violating**




$$K_{L,S} \rightarrow \mu^+ \mu^-$$

Dominated by Long-distance effects




$$K_L \rightarrow \pi^0 \nu \bar{\nu}$$

- Within the SM determined entirely from the weak effective Hamiltonian (**short-distance physics**)
- Purely **CP-violating**




$$K_{L,S} \rightarrow \mu^+ \mu^-$$

Dominated by Long-d. effects




$$K_L \rightarrow \pi^0 \nu \bar{\nu}$$

- Within the SM determined entirely from the weak effective Hamiltonian (*short-distance physics*)
- Purely **CP-violating**



$$K_{L,S} \rightarrow \mu^+ \mu^-$$

Exactly the same CKM dependence



In numbers

$$\mathcal{B}(K_L \rightarrow \mu^+ \mu^-) = (6.84 \pm 0.11) \cdot 10^{-9}$$

$$\mathcal{B}(K_S \rightarrow \mu^+ \mu^-)_{\text{LHCb}} < 2.1 \cdot 10^{-10}$$

LD SD

$$\mathcal{B}(K_L \rightarrow \mu^+ \mu^-)_{\text{SM}} = \begin{cases} (6.85 \pm 0.80 \pm 0.06) \times 10^{-9} (+) \\ (8.11 \pm 1.49 \pm 0.13) \times 10^{-9} (-) \end{cases}$$

LD SD

$$\mathcal{B}(K_S \rightarrow \mu^+ \mu^-)_{\text{SM}} = (5.18 \pm 1.50 \pm 0.02) \times 10^{-12}$$

How can we get rid of the
long-distance contributions?



The key: CPV comes from
the weak (SD) diagrams

Framework

We work under the following approximations –

1. We neglect CPV in mixing, negligible for our purposes, $\mathcal{O}(\varepsilon_K) \sim 10^{-3}$
2. We assume that the leptonic current is vectorial – within the SM this is an approximation, good to $\mathcal{O}(m_K^2/m_W^2)$
3. We neglect CPV in the long-distance physics, of $\mathcal{O}(\lambda^4) \sim 10^{-3}$

All fulfilled in the SM to $\mathcal{O}(10^{-3})$

CP analysis of $K \rightarrow \mu^+ \mu^-$

Initial state: kaon mass eigenstates are CP eigenstates,

$$\begin{array}{cc} K_L & K_S \\ \text{CP-odd} & \text{CP-even} \end{array}$$

Final state: since the kaon has $J = 0$, the dimuon state can have either $S = 0, \ell = 0$
corresponding to final states: or $S = 1, \ell = 1$

$$\begin{array}{cc} (\bar{\mu}\mu)_{\ell=0} & (\bar{\mu}\mu)_{\ell=1} \\ \text{CP-odd} & \text{CP-even} \end{array}$$

CP analysis of $K \rightarrow \mu^+ \mu^-$

Initial state: kaon mass eigenstates are CP eigenstates,

$$\begin{array}{cc} K_L & K_S \\ \text{CP-odd} & \text{CP-even} \end{array}$$

Final state: since the kaon has $J = 0$, the dimuon state can have either $S = 0, \ell = 0$
or $S = 1, \ell = 1$
corresponding to final states:

$$\begin{array}{cc} (\bar{\mu}\mu)_{\ell=0} & (\bar{\mu}\mu)_{\ell=1} \\ \text{CP-odd} & \text{CP-even} \end{array}$$

In practice, we measure the incoherent sums,

$$\Gamma(K_S \rightarrow \mu^+ \mu^-)_{meas.} = \Gamma(K_S \rightarrow (\mu^+ \mu^-)_{\ell=0}) + \Gamma(K_S \rightarrow (\mu^+ \mu^-)_{\ell=1})$$

$$\Gamma(K_L \rightarrow \mu^+ \mu^-)_{meas.} = \Gamma(K_L \rightarrow (\mu^+ \mu^-)_{\ell=0}) + \Gamma(K_L \rightarrow (\mu^+ \mu^-)_{\ell=1})$$

If we could extract the CPV modes, we would have a similar situation (theoretically) to $K_L \rightarrow \pi^0 \nu \bar{\nu}$

Time dependent rate

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

$$\left(\frac{d\Gamma}{dt}\right) = N_f f(t),$$

$$f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$

4 Experimental parameters

$$\{C_L, C_S, C_{Int.}, \varphi_0\}$$

Time dependent rate

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
 AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

$$\left(\frac{d\Gamma}{dt}\right) = N_f f(t), \quad f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$

4 Experimental parameters

Theory parameters: 4 amplitudes and 2 phases

$$\begin{array}{ll} \text{SD } |A(K_S \rightarrow (\mu^+ \mu^-)_{\ell=0})| & \\ |A(K_S \rightarrow (\mu^+ \mu^-)_{\ell=1})| & \varphi_0 \equiv \arg(A(K_S)_0^* A(K_L)_0) \\ |A(K_L \rightarrow (\mu^+ \mu^-)_{\ell=0})| & \varphi_1 \equiv \arg(A(K_S)_1^* A(K_L)_1) \\ \text{SD } |A(K_L \rightarrow (\mu^+ \mu^-)_{\ell=1})| & \end{array}$$

A priori, 6 theory parameters

Time dependent rate

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

$$\left(\frac{d\Gamma}{dt}\right) = N_f f(t), \quad f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$

4 Experimental parameters
 $\{C_L, C_S, C_{Int.}, \varphi_0\}$

Theory parameters: 4 amplitudes and 2 phases

$$\begin{aligned} \text{SD } & |A(K_S \rightarrow (\mu^+ \mu^-)_{\ell=0})| \\ & |A(K_S \rightarrow (\mu^+ \mu^-)_{\ell=1})| \\ & |A(K_L \rightarrow (\mu^+ \mu^-)_{\ell=0})| \\ \text{SD } & \cancel{|A(K_L \rightarrow (\mu^+ \mu^-)_{\ell=1})|} \end{aligned} \quad \begin{aligned} \varphi_0 &\equiv \arg(A(K_S)_0^* A(K_L)_0) \\ \varphi_1 &\equiv \arg(A(K_S)_1^* A(K_L)_1) \end{aligned}$$

Under our approximations two are negligible

A priori, 6 theory parameters

4, 1 of which is pure SD

$$\{|A(K_S)_0|, |A(K_L)_0|, |A(K_S)_1|, \arg(A(K_S)_0^* A(K_L)_0)\}$$

Entire system can be solved

4 Experimental parameters

$$\{C_L, C_S, C_{Int.}, \varphi_0\}$$



4 theory parameters

$$\{|A(K_S)_0|, |A(K_L)_0|, |A(K_S)_1|, \arg(A(K_S)_0^* A(K_L)_0)\}$$

$$C_L = |A(K_L)_0|^2$$

$$C_S = |A(K_S)_0|^2 + \beta_\mu^2 |A(K_S)_1|^2$$

$$C_{Int.} = D |A(K_S)_0 A(K_L)_0|$$

$$\varphi_0 = \arg(A(K_S)_0 A(K_L)_0)$$

$$D = \frac{N_{K^0} - N_{\bar{K}^0}}{N_{K^0} + N_{\bar{K}^0}}$$

In particular, we can solve for the CPV amplitude $|A(K_S)_0|$

$$\frac{1}{D^2} \frac{C_{Int.}^2}{C_L} = |A(K_S)_0|^2$$

SM prediction

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

$$|A(K_S)_0|^2 = \left| \frac{G_F}{2} \frac{2\alpha_{em} m_K m_\mu Y(x_t)}{\pi \sin^2 \theta_W} \times f_K \times V_{cs} V_{cd} \operatorname{Im} \left(\frac{V_{ts}^* V_{td}}{V_{cs}^* V_{cd}} \right) \right|^2$$

SM prediction

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

Only hadronic parameter, $\mathcal{O}(1\%)$
uncertainty from isospin breaking

$$|A(K_S)_0|^2 = \left| \frac{G_F}{2} \frac{2\alpha_{em} m_K m_\mu Y(x_t)}{\pi \sin^2 \theta_W} \times f_K \times V_{cs} V_{cd} \operatorname{Im} \left(\frac{V_{ts}^* V_{td}}{V_{cs}^* V_{cd}} \right) \right|^2$$

SM prediction

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

Only hadronic parameter, $\mathcal{O}(1\%)$
uncertainty from isospin breaking

$$|A(K_S)_0|^2 = \left| \frac{G_F}{2} \frac{2\alpha_{em} m_K m_\mu Y(x_t)}{\pi \sin^2 \theta_W} \times f_K \times V_{cs} V_{cd} \underbrace{\operatorname{Im} \left(\frac{V_{ts}^* V_{td}}{V_{cs}^* V_{cd}} \right)}_{\bar{\eta}_{ds}} \right|^2$$

$$\bar{\eta}_{ds} = A^2 \lambda^4 \bar{\eta} + \mathcal{O}(\lambda^7)$$


SM prediction

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

Only hadronic parameter, $\mathcal{O}(1\%)$
uncertainty from isospin breaking

$$|A(K_S)_0|^2 = \left| \frac{G_F}{2} \frac{2\alpha_{em} m_K m_\mu Y(x_t)}{\pi \sin^2 \theta_W} \times f_K \times V_{cs} V_{cd} \operatorname{Im} \left(\frac{V_{ts}^* V_{td}}{V_{cs}^* V_{cd}} \right) \right|^2$$

Current error on $\bar{\eta}_{ds}$
from B physics is $\mathcal{O}(5\%)$



$$\bar{\eta}_{ds} = A^2 \lambda^4 \bar{\eta} + \mathcal{O}(\lambda^7)$$

SM prediction

G. D'Ambrosio and T. Kitahara [arXiv:1707.06999]
AD, M. Ghosh, Y. Grossman, S. Schacht [arXiv:2104.06427]

Only hadronic parameter, $\mathcal{O}(1\%)$
uncertainty from isospin breaking

$$|A(K_S)_0|^2 = \left| \frac{G_F}{2} \frac{2\alpha_{em} m_K m_\mu Y(x_t)}{\pi \sin^2 \theta_W} \times f_K \times V_{cs} V_{cd} \operatorname{Im} \left(\frac{V_{ts}^* V_{td}}{V_{cs}^* V_{cd}} \right) \right|^2$$

$$\bar{\eta}_{ds} = A^2 \lambda^4 \bar{\eta} + \mathcal{O}(\lambda^7)$$

THE message for this talk:

A measurement of $\Gamma(K \rightarrow \mu^+ \mu^-)(t)$ would provide
a **clean** determination of $\bar{\eta}$ from **kaon physics**,
with ultimate theory uncertainty under 1 %

→ **golden mode**

Complementarity with $K_L \rightarrow \pi^0 \nu \bar{\nu}$

$$R_{\text{SL}} = \frac{\mathcal{B}(K_S \rightarrow \mu^+ \mu^-)_{\text{SD}}}{\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu})} = 1.55 \times 10^{-2} \left[\frac{\lambda}{0.225} \right]^2 \left[\frac{Y(x_t)}{X(x_t)} \right]^2$$

[A. J. Buras and E. Venturini,
[arXiv:2109.11032]]

Eliminates the dependence on $|V_{cb}|^4$

Complementary probe of NP models

[AD, M. Ghosh
[arXiv:2112.05801]]

How feasible is this experimentally?

Next talk by Rado Marchevski!


Some theory input on feasibility

AD, M. Ghosh, Y. Grossman, S. Schacht
(to appear)

What is the value of φ_0 ?

Will affect the sensitivity of the measurement

$$f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$




Some theory input on feasibility

AD, M. Ghosh, Y. Grossman, S. Schacht
(to appear)

What is the value of φ_0 ?

$$f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$

Will affect the sensitivity of the measurement

$$\cos^2 \varphi_0 = \frac{(\text{Im} A(K_L))^2}{|A(K_L)|^2} = \frac{\text{Diagram}}{\mathcal{B}(K_L \rightarrow \mu^+ \mu^-)} = C_{\text{QED}} \frac{\mathcal{B}(K_L \rightarrow \gamma\gamma)}{\mathcal{B}(K_L \rightarrow \mu^+ \mu^-)}$$



Some theory input on feasibility

AD, M. Ghosh, Y. Grossman, S. Schacht
(to appear)

What is the value of φ_0 ?

$$f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$

Will affect the sensitivity of the measurement


$$\cos^2 \varphi_0 = \frac{(\text{Im} A(K_L))^2}{|A(K_L)|^2} = \frac{\text{Diagram}}{\mathcal{B}(K_L \rightarrow \mu^+ \mu^-)} = C_{\text{QED}} \frac{\mathcal{B}(K_L \rightarrow \gamma\gamma)}{\mathcal{B}(K_L \rightarrow \mu^+ \mu^-)}$$


$$\left[\cos \varphi_0 \right]_{\text{SM}} = 0.978 \pm 0.009 \quad \text{Close to maximal}$$

Some theory input on feasibility

AD, M. Ghosh, Y. Grossman, S. Schacht
(to appear)

$$f(t) = C_L e^{-\Gamma_L t} + C_S e^{-\Gamma_S t} + 2 C_{Int.} \cos(\Delta m t - \varphi_0) e^{-\Gamma t},$$



Approximate SM
prediction,
 $D = 1$:

$$(C_L)_{SM} \equiv 1$$

$$(C_S)_{SM} \approx 0.43$$

$$(C_{Int.})_{SM} \approx 0.12$$

Naive estimate - $\mathcal{O}(10^{13})$ Kaons are needed

Summary and Conclusions

- ◆ The time-dependent rate in $K \rightarrow \mu^+ \mu^-$ is a complementary *golden mode*, allowing for a clean independent determination of CKM parameters from kaon physics.
- ◆ The ultimate theoretical uncertainty is of $\mathcal{O}(1\%)$
- ◆ The relevant CKM parameter is $\bar{\eta}_{ds} \approx A^2 \lambda^4 \bar{\eta}$, the same as for $K_L \rightarrow \pi^0 \bar{\nu} \nu$.
- ◆ The ratio $\mathcal{B}(K_S \rightarrow \mu^+ \mu^-)_{\ell=0} / \mathcal{B}(K_L \rightarrow \pi^0 \bar{\nu} \nu)$ is an extremely clean SM observable, dependent only on the parameters λ , m_t .
[A. J. Buras and E. Venturini, [arXiv:2109.11032]]
- ◆ The same $K \rightarrow \mu^+ \mu^-$ observable is also a *sensitive probe of NP*
[AD, M. Ghosh [arXiv:2112.05801]]
- ◆ The phase shift in the oscillating rate, φ_0 , is an additional clean SM prediction, and implies close to maximal integrated event rate.
[AD, Ghosh, Grossman, Schacht [in preparation]]
- ◆ A theorist's estimation - we need $\mathcal{O}(10^{13})$ neutral kaons, with $\mathcal{O}(1)$ Dilution factor

Thank you