

KEK-PH 2023


# $S_4$ flavor model with 3HDM

Hiroshima Univ. Yukimura Izawa

November 8, 2023



Graduate School of  
Advanced Science and Engineering  
Hiroshima University



**SKCM<sup>2</sup>**  
WPI HIROSHIMA UNIVERSITY

## Contents

- 1 . Introduction
- 2 .  $S_4$  symmetry
- 3 . 3 HDM
- 4 . Flavor model
- 5 . Numerical calculation
- 6 . Potential analysis
- 7 . Summary

# 1. Introduction

## Fundamental forces

(strong, weak, electromagnetic and gravitational interactions)

Standard model(SM)

$SU(3) \times SU(2)_L \times U(1)_Y$  gauge symmetry

Quarks and leptons (SM particles)

→ generation structure

(mass differences and flavor mixing)

Especially leptons → large flavor mixing



It can't be explained in SM. (SM only assign parameter.)

Standard Model of Elementary Particles and Gravity

three generations of matter (fermions)			interactions / force carriers (bosons)		
	I	II	III		
mass	$\approx 2.2 \text{ MeV}/c^2$	$\approx 1.28 \text{ GeV}/c^2$	$\approx 173.1 \text{ GeV}/c^2$	0	$\approx 124.97 \text{ GeV}/c^2$
charge	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0	0
spin	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	0
	<b>u</b> up	<b>c</b> charm	<b>t</b> top	<b>g</b> gluon	<b>H</b> higgs
	<b>d</b> down	<b>s</b> strange	<b>b</b> bottom	<b>γ</b> photon	<b>G</b> graviton
	<b>e</b> electron	<b>μ</b> muon	<b>τ</b> tau	<b>Z</b> Z boson	
	<b>ν<sub>e</sub></b> electron neutrino	<b>ν<sub>μ</sub></b> muon neutrino	<b>ν<sub>τ</sub></b> tau neutrino	<b>W</b> W boson	

QUARKS (I, II, III)  
 LEPTONS (e, μ, τ, ν<sub>e</sub>, ν<sub>μ</sub>, ν<sub>τ</sub>)  
 GAUGE BOSONS VECTOR BOSONS (g, γ, Z, W)  
 SCALAR BOSONS (H)  
 HYPOTHETICAL TENSOR BOSONS (G)

<https://www.wikiwand.com/>

# 1. Introduction

Altareli and Feruglio impose discrete symmetry(flavor symmetry) among generations.

G. Altarelli and F. Feruglio, Nucl. Phys. B741 (2006), 215–235.



In this study, we choose  $S_4$  symmetry as flavor symmetry.

In addition, we suppose three Higgs doublets model(3HDM).

We built new flavor model and perform the analysis.

Standard Model of Elementary Particles and Gravity

three generations of matter (fermions)			interactions / force carriers (bosons)	
I	II	III		
mass $\approx 2.2 \text{ MeV}/c^2$ charge $\frac{2}{3}$ spin $\frac{1}{2}$ <b>u</b> up	mass $\approx 1.28 \text{ GeV}/c^2$ charge $\frac{2}{3}$ spin $\frac{1}{2}$ <b>c</b> charm	mass $\approx 173.1 \text{ GeV}/c^2$ charge $\frac{2}{3}$ spin $\frac{1}{2}$ <b>t</b> top	mass 0 charge 0 spin 1 <b>g</b> gluon	mass $\approx 124.97 \text{ GeV}/c^2$ charge 0 spin 0 <b>H</b> higgs
mass $\approx 4.7 \text{ MeV}/c^2$ charge $-\frac{1}{3}$ spin $\frac{1}{2}$ <b>d</b> down	mass $\approx 96 \text{ MeV}/c^2$ charge $-\frac{1}{3}$ spin $\frac{1}{2}$ <b>s</b> strange	mass $\approx 4.18 \text{ GeV}/c^2$ charge $-\frac{1}{3}$ spin $\frac{1}{2}$ <b>b</b> bottom	mass 0 charge 0 spin 1 <b><math>\gamma</math></b> photon	mass 0 charge 0 spin 2 <b>G</b> graviton
mass $\approx 0.511 \text{ MeV}/c^2$ charge -1 spin $\frac{1}{2}$ <b>e</b> electron	mass $\approx 105.66 \text{ MeV}/c^2$ charge -1 spin $\frac{1}{2}$ <b><math>\mu</math></b> muon	mass $\approx 1.7768 \text{ GeV}/c^2$ charge -1 spin $\frac{1}{2}$ <b><math>\tau</math></b> tau	mass $\approx 91.19 \text{ GeV}/c^2$ charge 0 spin 1 <b>Z</b> Z boson	
mass $< 1.0 \text{ eV}/c^2$ charge 0 spin $\frac{1}{2}$ <b><math>\nu_e</math></b> electron neutrino	mass $< 0.17 \text{ MeV}/c^2$ charge 0 spin $\frac{1}{2}$ <b><math>\nu_\mu</math></b> muon neutrino	mass $< 18.2 \text{ MeV}/c^2$ charge 0 spin $\frac{1}{2}$ <b><math>\nu_\tau</math></b> tau neutrino	mass $\approx 80.39 \text{ GeV}/c^2$ charge $\pm 1$ spin 1 <b>W</b> W boson	

QUARKS (left column), LEPTONS (right column), GAUGE BOSONS VECTOR BOSONS (middle), SCALAR BOSONS (right), HYPOTHETICAL TENSOR BOSONS (far right).

<https://www.wikiwand.com/>

## 2. $S_4$ symmetry


$S_4$  symmetry : Fourth order symmetric group

$$(x_1, x_2, x_3, x_4) \rightarrow (x_i, x_j, x_k, x_l) \quad 4! = 24 \text{ elements}$$

Representation  $1, 1', 2, 3, 3'$

$A_4$  symmetry : Fourth order alternating group

Representation  $1, 1', 1'', 3_S, 3_A$  12 elements



Multiplication rule

$$3 \times 3 = 1 + 2 + 3 + 3'$$

$$3' \times 3' = 1 + 2 + 3 + 3'$$

$$3 \times 3' = 1' + 2 + 3 + 3'$$

$$2 \times 2 = 1 + 1' + 2$$

$$2 \times 3 = 3 + 3'$$

$$2 \times 3' = 3 + 3'$$

$$3 \times 1' = 3'$$

$$3' \times 1' = 3$$

$$2 \times 1' = 2$$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}_2 \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}_2 = (\alpha_1\beta_1 + \alpha_2\beta_2)_1 \oplus (-\alpha_1\beta_2 + \alpha_2\beta_1)_{1'} \oplus \begin{pmatrix} \alpha_1\beta_2 + \alpha_2\beta_1 \\ \alpha_1\beta_1 - \alpha_2\beta_2 \end{pmatrix}_2$$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix}_3 \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}_3 = (\alpha_1\beta_1 + \alpha_2\beta_2 + \alpha_3\beta_3)_1 \oplus \begin{pmatrix} \frac{1}{\sqrt{2}}(\alpha_2\beta_2 - \alpha_3\beta_3) \\ \frac{1}{\sqrt{6}}(-2\alpha_1\beta_1 + \alpha_2\beta_2 + \alpha_3\beta_3) \end{pmatrix}_2$$

$$\oplus \begin{pmatrix} \alpha_3\beta_2 + \alpha_2\beta_3 \\ \alpha_1\beta_3 + \alpha_3\beta_1 \\ \alpha_2\beta_1 + \alpha_1\beta_2 \end{pmatrix}_3 \oplus \begin{pmatrix} \alpha_3\beta_2 - \alpha_2\beta_3 \\ \alpha_1\beta_3 - \alpha_3\beta_1 \\ \alpha_2\beta_1 - \alpha_1\beta_2 \end{pmatrix}_{3'}$$

### 3. 3HDM

Extend SM Higgs doublet to 3 (12 real scalar fields)

$$\phi_1 = \begin{pmatrix} \phi_1^+ \\ \phi_1^0 \end{pmatrix}, \phi_2 = \begin{pmatrix} \phi_2^+ \\ \phi_2^0 \end{pmatrix}, \phi_3 = \begin{pmatrix} \phi_3^+ \\ \phi_3^0 \end{pmatrix}$$

Higgs Potential in 3HDM under  $SU(2)_L \otimes U(1)_Y$

$$V = - \sum_{i,j=1}^3 m_{ij}^2 (\phi_i^\dagger \phi_j) + \frac{1}{2} \sum_{i,j,k,l=1}^3 \lambda_{ijkl} (\phi_i^\dagger \phi_j) (\phi_k^\dagger \phi_l)$$

Potential minimum conditions

$$\left( \frac{\partial V}{\partial \phi_1} \right)_{\phi_1=\langle \phi_1 \rangle, \phi_2=\langle \phi_2 \rangle, \phi_3=\langle \phi_3 \rangle} = 0$$

$$\left( \frac{\partial V}{\partial \phi_2} \right)_{\phi_1=\langle \phi_1 \rangle, \phi_2=\langle \phi_2 \rangle, \phi_3=\langle \phi_3 \rangle} = 0$$

$$\left( \frac{\partial V}{\partial \phi_3} \right)_{\phi_1=\langle \phi_1 \rangle, \phi_2=\langle \phi_2 \rangle, \phi_3=\langle \phi_3 \rangle} = 0$$



Spontaneous symmetry breaking

3 degrees of freedom are eaten by  $W$  and  $Z$  bosons.

→  $\phi$  is represented by the expansion of 9 (=12-3) real scalar fields

$$\phi_i = \begin{pmatrix} \rho_i^+ \\ \frac{1}{\sqrt{2}}(v_i + \rho_i + i\chi_i) \end{pmatrix}, i = 1, 2, 3$$

mass eigenstates



- (i) Three CP-even scalar fields
- (ii) Two CP-odd scalar fields
- (iii) Four charged scalar fields

## 4. Flavor model

	$\bar{l} = (\bar{l}_e, \bar{l}_\mu, \bar{l}_\tau)$	$l_R = (e_R, \mu_R)$	$\tau_R$	$\nu_{eR}$	$\nu_R = (\nu_{\mu R}, \nu_{\tau R})$	$\phi = (\phi_1, \phi_2, \phi_3)$	$X$	$\Theta$
$SU(2)_L$	2	1	1	1	1	2	1	1
$S_4$	3	2	1	1	2	3	2	1
$U(1)_{FN}$	0	+1	0	0	0	0	-1	-1

Lagrangian :  $-L_Y = L_l + L_D + L_M + h.c.$

(1) Mass terms of charged leptons :  $L_l = \frac{y_{e\mu}}{\Lambda} \bar{l} \phi l_R \Theta + y_\tau \bar{l} \phi \tau_R + \frac{y_l}{\Lambda} \bar{l} \phi l_R X$

(2) Mass term of Dirac neutrino :  $L_D = y_{De} \bar{l} \tilde{\phi} \nu_{eR} + y_{D\mu\tau} \bar{l} \tilde{\phi} \nu_R$

(3) Mass term of right-handed Majorana neutrino :  $L_M = \frac{1}{2} M_{eR} \bar{\nu}_{eR}^c \nu_{eR} + \frac{1}{2} M_{\mu\tau R} \bar{\nu}_R^c \nu_R$



Calculate mass matrices of charged leptons and left-handed Majorana neutrino

## Calculation of mass matrices

### (1) Mass terms of charged leptons

$$L_l = \frac{y_{e\mu}}{\Lambda} \bar{l} \phi l_R \Theta + y_\tau \bar{l} \phi \tau_R + \frac{y_l}{\Lambda} \bar{l} \phi l_R X$$

$$\begin{aligned} \frac{y_l}{\Lambda} \begin{pmatrix} \bar{l}_e \\ \bar{l}_\mu \\ \bar{l}_\tau \end{pmatrix}_3 \otimes \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3 \otimes \begin{pmatrix} e_R \\ \mu_R \end{pmatrix}_2 \Theta &= \frac{y_{e\mu}}{\Lambda} \begin{pmatrix} \frac{1}{\sqrt{2}} (\bar{l}_\mu \phi_2 - \bar{l}_\tau \phi_3) \\ \frac{1}{\sqrt{6}} (-2\bar{l}_e \phi_1 + \bar{l}_\mu \phi_2 + \bar{l}_\tau \phi_3) \end{pmatrix}_2 \otimes \begin{pmatrix} e_R \\ \mu_R \end{pmatrix}_2 \Theta \\ &= \frac{y_{e\mu}}{\Lambda} \frac{1}{\sqrt{2}} (\bar{l}_\mu \phi_2 - \bar{l}_\tau \phi_3) e_R \Theta + \frac{y_{e\mu}}{\Lambda} \frac{1}{\sqrt{6}} (-2\bar{l}_e \phi_1 + \bar{l}_\mu \phi_2 + \bar{l}_\tau \phi_3) \mu_R \Theta \\ &\quad \Downarrow \quad \langle \Theta \rangle = \Theta_0, \langle \phi \rangle = (v_1, v_2, v_3) \\ &= \frac{y_{e\mu} \Theta_0}{\Lambda} \frac{1}{\sqrt{2}} (\bar{\mu}_L v_2 - \bar{\tau}_L v_3) e_R + \frac{y_{e\mu} \Theta_0}{\Lambda} \frac{1}{\sqrt{6}} (-2\bar{e}_L v_1 + \bar{\mu}_L v_2 + \bar{\tau}_L v_3) \mu_R \end{aligned}$$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix}_3 \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}_3 = \begin{pmatrix} \frac{1}{\sqrt{2}} (\alpha_2 \beta_2 - \alpha_3 \beta_3) \\ \frac{1}{\sqrt{6}} (-2\alpha_1 \beta_1 + \alpha_2 \beta_2 + \alpha_3 \beta_3) \end{pmatrix}_2$$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}_2 \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}_2 = (\alpha_1 \beta_1 + \alpha_2 \beta_2)_1$$




## Calculation of mass matrices

### (1) Mass terms of charged leptons

$$L_l = \frac{y_{e\mu}}{\Lambda} \bar{l} \phi l_R \Theta + \boxed{y_\tau \bar{l} \phi \tau_R} + \frac{y_l}{\Lambda} \bar{l} \phi l_R X$$

$$y_\tau \underbrace{\begin{pmatrix} \bar{l}_e \\ \bar{l}_\mu \\ \bar{l}_\tau \end{pmatrix}_3}_{1} \otimes \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3 \otimes \tau_R = y_\tau (\bar{l}_e \phi_1 + \bar{l}_\mu \phi_2 + \bar{l}_\tau \phi_3) \tau_R$$


 $\langle \phi \rangle = (v_1, v_2, v_3)$

$$= y_\tau (\bar{e}_L v_1 \tau_R + \bar{\mu}_L v_2 \tau_R + \bar{\tau}_L v_3 \tau_R)$$

Mass matrix of charged leptons  
consisting of one and two terms

$$M_{l1} = \begin{pmatrix} 0 & -\frac{2y_{e\mu}\Theta_0}{\sqrt{6}\Lambda} v_1 & y_\tau v_1 \\ \frac{y_{e\mu}\Theta_0}{\sqrt{2}\Lambda} v_2 & \frac{y_{e\mu}\Theta_0}{\sqrt{6}\Lambda} v_2 & y_\tau v_2 \\ -\frac{y_{e\mu}\Theta_0}{\sqrt{2}\Lambda} v_3 & \frac{y_{e\mu}\Theta_0}{\sqrt{6}\Lambda} v_3 & y_\tau v_3 \end{pmatrix}_{LR}$$

## Calculation of mass matrices

### (1) Mass terms of charged leptons

$$L_l = \frac{y_{e\mu}}{\Lambda} \bar{l}\phi l_R \Theta + y_\tau \bar{l}\phi \tau_R + \frac{y_l}{\Lambda} \bar{l}\phi l_R X$$

$$\begin{aligned} & \frac{y_l}{\Lambda} \underbrace{\begin{pmatrix} \bar{l}_e \\ \bar{l}_\mu \\ \bar{l}_\tau \end{pmatrix}_3}_{1,2} \otimes \underbrace{\begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3}_{1,2} \otimes \begin{pmatrix} e_R \\ \mu_R \end{pmatrix}_2 \otimes \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}_2 \\ &= \frac{y_{l1}}{\Lambda} (\bar{l}_e \phi_1 + \bar{l}_\mu \phi_2 + \bar{l}_\tau \phi_3)_1 \otimes (e_R X_1 + \mu_R X_2)_1 + \frac{y_{l2}}{\Lambda} \begin{pmatrix} \frac{1}{\sqrt{2}}(\bar{l}_\mu \phi_2 - \bar{l}_\tau \phi_3) \\ \frac{1}{\sqrt{6}}(-2\bar{l}_e \phi_1 + \bar{l}_\mu \phi_2 + \bar{l}_\tau \phi_3) \end{pmatrix}_2 \otimes \begin{pmatrix} e_R X_2 + \mu_R X_1 \\ e_R X_1 - \mu_R X_2 \end{pmatrix}_2 \\ &= \frac{y_{l1}}{\Lambda} (\bar{l}_e \phi_1 e_R X_1 + \bar{l}_e \phi_1 \mu_R X_2 + \bar{l}_\mu \phi_2 e_R X_1 + \bar{l}_\mu \phi_2 \mu_R X_2 + \bar{l}_\tau \phi_3 e_R X_1 + \bar{l}_\tau \phi_3 \mu_R X_2) \\ &+ \frac{y_{l2}}{\Lambda} \left[ \frac{1}{\sqrt{2}} (\bar{l}_\mu \phi_2 e_R X_2 + \bar{l}_\mu \phi_2 \mu_R X_1 - \bar{l}_\tau \phi_3 e_R X_2 - \bar{l}_\tau \phi_3 \mu_R X_1) \right. \\ &+ \left. \frac{1}{\sqrt{6}} (-2\bar{l}_e \phi_1 e_R X_1 + 2\bar{l}_e \phi_1 \mu_R X_2 + \bar{l}_\mu \phi_2 e_R X_1 - \bar{l}_\mu \phi_2 \mu_R X_2 + \bar{l}_\tau \phi_3 e_R X_1 - \bar{l}_\tau \phi_3 \mu_R X_2) \right] \end{aligned}$$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix}_3 \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}_3 = \begin{pmatrix} \frac{1}{\sqrt{2}}(\alpha_2 \beta_2 - \alpha_3 \beta_3) \\ \frac{1}{\sqrt{6}}(-2\alpha_1 \beta_1 + \alpha_2 \beta_2 + \alpha_3 \beta_3) \end{pmatrix}_2$$

$$\begin{aligned} \begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}_2 \otimes \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}_2 &= (\alpha_1 \beta_1 + \alpha_2 \beta_2)_1 \\ &\oplus (\alpha_1 \beta_2 + \alpha_2 \beta_1)_2 \\ &\oplus (\alpha_1 \beta_1 - \alpha_2 \beta_2)_2 \end{aligned}$$

## Calculation of mass matrices

### (1) Mass terms of charged leptons

$$L_l = \frac{y_{e\mu}}{\Lambda} \bar{l}\phi l_R \Theta + y_\tau \bar{l}\phi \tau_R + \frac{y_l}{\Lambda} \bar{l}\phi l_R X$$

$$\langle \phi \rangle = (v_1, v_2, v_3), \langle X \rangle = (X_1, 0)$$

Mass matrix of charged leptons  
consisting of third term

$$M_{l2} = \begin{pmatrix} \left(\frac{y_{l1}}{\Lambda} - \frac{2y_{l2}}{\sqrt{6}\Lambda}\right) v_1 X_1 & 0 & 0 \\ \left(\frac{y_{l1}}{\Lambda} + \frac{y_{l2}}{\sqrt{6}\Lambda}\right) v_2 X_1 & \frac{y_{l2}}{\sqrt{2}\Lambda} v_2 X_1 & 0 \\ \left(\frac{y_{l1}}{\Lambda} - \frac{y_{l2}}{\sqrt{6}\Lambda}\right) v_3 X_1 & -\frac{y_{l2}}{\sqrt{2}\Lambda} v_3 X_1 & 0 \end{pmatrix}_{LR}$$

### Mass matrix of charged leptons

$$M_l = M_{l1} + M_{l2} = \begin{pmatrix} 0 & -\frac{2y_{e\mu}\Theta_0}{\sqrt{6}\Lambda} v_1 & y_\tau v_1 \\ \frac{y_{e\mu}\Theta_0}{\sqrt{2}\Lambda} v_2 & \frac{y_{e\mu}\Theta_0}{\sqrt{6}\Lambda} v_2 & y_\tau v_2 \\ -\frac{y_{e\mu}\Theta_0}{\sqrt{2}\Lambda} v_3 & \frac{y_{e\mu}\Theta_0}{\sqrt{6}\Lambda} v_3 & y_\tau v_3 \end{pmatrix}_{LR} + \begin{pmatrix} \left(\frac{y_{l1}}{\Lambda} - \frac{2y_{l2}}{\sqrt{6}\Lambda}\right) v_1 X_1 & 0 & 0 \\ \left(\frac{y_{l1}}{\Lambda} + \frac{y_{l2}}{\sqrt{6}\Lambda}\right) v_2 X_1 & \frac{y_{l2}}{\sqrt{2}\Lambda} v_2 X_1 & 0 \\ \left(\frac{y_{l1}}{\Lambda} - \frac{y_{l2}}{\sqrt{6}\Lambda}\right) v_3 X_1 & -\frac{y_{l2}}{\sqrt{2}\Lambda} v_3 X_1 & 0 \end{pmatrix}_{LR}$$

## Calculation of mass matrices

(2) Mass terms of Dirac neutrino

$$L_D = y_{De} \bar{l} \tilde{\phi} \nu_{eR} + y_{D\mu\tau} \bar{l} \tilde{\phi} \nu_R \quad \text{Mass matrix of Dirac neutrino} \quad M_D = \begin{pmatrix} y_{De} v_1 & 0 & -2/\sqrt{6} y_{D\mu\tau} v_1 \\ y_{De} v_2 & 1/\sqrt{2} y_{D\mu\tau} v_2 & 1/\sqrt{6} y_{D\mu\tau} v_2 \\ y_{De} v_3 & -1/\sqrt{2} y_{D\mu\tau} v_3 & 1/\sqrt{6} y_{D\mu\tau} v_3 \end{pmatrix}_{LR}$$

(3) Mass terms of Majorana neutrino

$$L_M = \frac{1}{2} M_{eR} \bar{\nu}_{eR}^c \nu_{eR} + \frac{1}{2} M_{\mu\tau R} \bar{\nu}_R^c \nu_R \quad \text{Mass matrix of Majorana neutrino} \quad M_R = \begin{pmatrix} M_{eR} & 0 & 0 \\ 0 & M_{\mu\tau R} & 0 \\ 0 & 0 & M_{\mu\tau R} \end{pmatrix}$$

By using type-1 seesaw mechanism, Mass matrix of neutrino

$$m_\nu = -M_D M_R^{-1} M_D^T$$

$$m_\nu = \begin{pmatrix} -\frac{e^{2i\phi} y_{De} y_{De}^2 v_1^2}{M_{eR}} - \frac{2e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_1^2}{3M_{\mu\tau R}} & -\frac{e^{2i\phi} y_{De} y_{De}^2 v_1 v_2}{M_{eR}} + \frac{e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_1 v_2}{3M_{\mu\tau R}} & -\frac{e^{2i\phi} y_{De} y_{De}^2 v_3 v_1}{M_{eR}} + \frac{e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_3 v_1}{3M_{\mu\tau R}} \\ -\frac{e^{2i\phi} y_{De} y_{De}^2 v_1 v_2}{M_{eR}} + \frac{e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_1 v_2}{3M_{\mu\tau R}} & -\frac{e^{2i\phi} y_{De} y_{De}^2 v_2^2}{M_{eR}} - \frac{2e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_2^2}{3M_{\mu\tau R}} & -\frac{e^{2i\phi} y_{De} y_{De}^2 v_2 v_3}{M_{eR}} + \frac{e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_2 v_3}{3M_{\mu\tau R}} \\ -\frac{e^{2i\phi} y_{De} y_{De}^2 v_3 v_1}{M_{eR}} + \frac{e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_3 v_1}{3M_{\mu\tau R}} & -\frac{e^{2i\phi} y_{De} y_{De}^2 v_2 v_3}{M_{eR}} + \frac{e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_2 v_3}{3M_{\mu\tau R}} & -\frac{e^{2i\phi} y_{De} y_{De}^2 v_3^2}{M_{eR}} - \frac{2e^{2i\phi} y_{\mu\tau} y_{D\mu\tau}^2 v_3^2}{3M_{\mu\tau R}} \end{pmatrix}$$


## 5. Numerical calculation

Satisfy  $m_e, m_\mu, m_\tau, \Delta m_{21}^2, \Delta m_{31}^2, \sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13}$

Parameter  $\alpha, \beta, y_{e\mu}, y_\tau, y_{l1}, y_{l2}, X_1, m_1, y_{De}, y_{D\mu\tau}, \phi_{yDe}$


Prediction  $\sin^2 \theta_{12}, \sin^2 \theta_{23}, \sin^2 \theta_{13}, \delta_{CP}, m_{light}, m_1 + m_2 + m_3,$   
 $m_{ee}, \eta_1, \eta_2$

The relation between  $\alpha$  and  $\beta$




$$\begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} v \sin \alpha \\ v \cos \alpha \sin \beta \\ v \cos \alpha \cos \beta \end{pmatrix}$$

The relation between  $v_1$  and  $v_2$




The relation between  $v_1$  and  $v_3$



## Numerical calculation

Prediction of  $\delta_{CP}$  and  $\sin^2\theta_{23}$



NuFIT 5.2  
 $0.408 \leq \sin^2\theta_{23} \leq 0.603$


$0.488 \leq \sin^2\theta_{23} \leq 0.603$

$\delta_{CP} \approx -67.7^\circ$

Strong prediction of  $\delta_{CP}$


## Numerical calculation

Prediction of the effective neutrino mass  $m_{ee}$  in the  $0\nu\beta\beta$  decay experiment and the lightest neutrino mass  $m_{\text{light}}$



Our model can be confirmed in the near future.

$0\nu\beta\beta$  decay



[https://en.wikipedia.org/wiki/Double\\_beta\\_decay](https://en.wikipedia.org/wiki/Double_beta_decay)

Decay rate


$$\Gamma \propto m_{ee}^2$$

Effective mass of electron neutrino

$$m_{ee} = |m_1 U_{e1}^2 + m_2 U_{e2}^2 + m_3 U_{e3}^2|$$

## Numerical calculation

Prediction of sum of neutrino mass  
 $m_1 + m_2 + m_3$  and  $\delta_{CP}$




Limit from cosmology  
(arXiv:1807.06209)  
 $m_1 + m_2 + m_3 \leq 0.12\text{eV}$



## Numerical calculation

Prediction of Majorana phases  $\eta_1, \eta_2$



$$\eta_1 = \arg \left[ \frac{U_{e1} U_{e3}^*}{\cos\theta_{12} \cos\theta_{13} \sin\theta_{13} e^{i\delta_{CP}}} \right], \quad \eta_2 = \arg \left[ \frac{U_{e2} U_{e3}^*}{\sin\theta_{12} \cos\theta_{13} \sin\theta_{13} e^{i\delta_{CP}}} \right]$$

## 6. Potential analysis

Check the Higgs VEV  $\langle \phi \rangle = (v_1, v_2, v_3)$

### 3HDM + $S_4$ symmetry

Consider  $\phi$  as  $S_4$  triplet  $\phi = (\phi_1, \phi_2, \phi_3)$

Calculate Higgs potential  $V = -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 + c \phi^\dagger \phi X^\dagger X + k \phi^\dagger \phi \Theta^\dagger \Theta + (g \phi^\dagger \phi X^\dagger \Theta + h.c.)$

$$\phi^\dagger \phi = \begin{pmatrix} \phi_1^\dagger \\ \phi_2^\dagger \\ \phi_3^\dagger \end{pmatrix}_3 \otimes \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3 = (\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 + \phi_3^\dagger \phi_3)_1 = |\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2$$

## Potential analysis

Calculate Higgs potential  $V = -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 + c \phi^\dagger \phi X^\dagger X + k \phi^\dagger \phi \Theta^\dagger \Theta + (g \phi^\dagger \phi \Theta^\dagger X + h.c.)$

$$\begin{aligned}
 (\phi^\dagger \phi)^2 &= \underbrace{\begin{pmatrix} \phi_1^\dagger \\ \phi_2^\dagger \\ \phi_3^\dagger \end{pmatrix}_3}_{1,2,3,3'} \otimes \underbrace{\begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3}_{1,2,3,3'} = (|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2)^2 \oplus \underbrace{\begin{pmatrix} \frac{1}{\sqrt{2}}(\phi_2^\dagger \phi_2 - \phi_3^\dagger \phi_3) \\ \frac{1}{\sqrt{6}}(-2\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 + \phi_3^\dagger \phi_3) \end{pmatrix}_2}_{1,2,3,3'} \otimes \underbrace{\begin{pmatrix} \frac{1}{\sqrt{2}}(\phi_2^\dagger \phi_2 - \phi_3^\dagger \phi_3) \\ \frac{1}{\sqrt{6}}(-2\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 + \phi_3^\dagger \phi_3) \end{pmatrix}_2}_{1,2,3,3'} \\
 &\oplus \underbrace{\begin{pmatrix} \phi_3^\dagger \phi_2 + \phi_2^\dagger \phi_3 \\ \phi_1^\dagger \phi_3 + \phi_3^\dagger \phi_1 \\ \phi_2^\dagger \phi_1 + \phi_1^\dagger \phi_2 \end{pmatrix}_3}_{1,2,3,3'} \otimes \underbrace{\begin{pmatrix} \phi_3^\dagger \phi_2 + \phi_2^\dagger \phi_3 \\ \phi_1^\dagger \phi_3 + \phi_3^\dagger \phi_1 \\ \phi_2^\dagger \phi_1 + \phi_1^\dagger \phi_2 \end{pmatrix}_3}_{1,2,3,3'} \oplus \underbrace{\begin{pmatrix} \phi_3^\dagger \phi_2 - \phi_2^\dagger \phi_3 \\ \phi_1^\dagger \phi_3 - \phi_3^\dagger \phi_1 \\ \phi_2^\dagger \phi_1 - \phi_1^\dagger \phi_2 \end{pmatrix}_{3'}}_{1,2,3,3'} \otimes \underbrace{\begin{pmatrix} \phi_3^\dagger \phi_2 - \phi_2^\dagger \phi_3 \\ \phi_1^\dagger \phi_3 - \phi_3^\dagger \phi_1 \\ \phi_2^\dagger \phi_1 - \phi_1^\dagger \phi_2 \end{pmatrix}_{3'}}_{1,2,3,3'} \\
 &= (|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2)^2 \\
 &\oplus \frac{2}{3} (|\phi_1|^4 + |\phi_2|^4 + |\phi_3|^4 - |\phi_1|^2 |\phi_2|^2 - |\phi_2|^2 |\phi_3|^2 - |\phi_3|^2 |\phi_1|^2) \\
 &\oplus \left[ (\phi_1^\dagger \phi_2)^2 + (\phi_2^\dagger \phi_3)^2 + (\phi_3^\dagger \phi_1)^2 + |\phi_1|^2 |\phi_2|^2 + |\phi_2|^2 |\phi_3|^2 + |\phi_3|^2 |\phi_1|^2 + h.c. \right] \\
 &\oplus \left[ (\phi_1^\dagger \phi_2)^2 + (\phi_2^\dagger \phi_3)^2 + (\phi_3^\dagger \phi_1)^2 - |\phi_1|^2 |\phi_2|^2 - |\phi_2|^2 |\phi_3|^2 - |\phi_3|^2 |\phi_1|^2 + h.c. \right]
 \end{aligned}$$

## Potential analysis

Calculate Higgs potential

$$V = -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 + c \phi^\dagger \phi X^\dagger X + k \phi^\dagger \phi \Theta^\dagger \Theta + (g \phi^\dagger \phi \Theta^\dagger X + h.c.)$$

$$\phi^\dagger \phi X^\dagger X = \underbrace{\begin{pmatrix} \phi_1^\dagger \\ \phi_2^\dagger \\ \phi_3^\dagger \end{pmatrix}_3 \otimes \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3}_{1,2} \otimes \underbrace{\begin{pmatrix} X_1^\dagger \\ X_2^\dagger \end{pmatrix}_2 \otimes \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}_2}_{1,2} = (\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 + \phi_3^\dagger \phi_3)_1 \otimes (X_1^\dagger X_1 + X_2^\dagger X_2)_1 + \begin{pmatrix} \frac{1}{\sqrt{2}}(\phi_2^\dagger \phi_2 - \phi_3^\dagger \phi_3) \\ \frac{1}{\sqrt{6}}(-2\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 + \phi_3^\dagger \phi_3) \end{pmatrix}_2 \otimes \begin{pmatrix} X_1^\dagger X_2 + X_2^\dagger X_1 \\ X_1^\dagger X_1 - X_2^\dagger X_2 \end{pmatrix}_2$$

$\langle X \rangle = (X_1, 0)$

$$= |X_1|^2 (|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2) + \frac{|X_1|^2}{\sqrt{6}} (-2|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2)$$

$$\phi^\dagger \phi \Theta^\dagger \Theta = \begin{pmatrix} \phi_1^\dagger \\ \phi_2^\dagger \\ \phi_3^\dagger \end{pmatrix}_3 \otimes \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3 \Theta^\dagger \Theta = |\Theta_0|^2 (|\phi_1|^2 + |\phi_2|^2 + |\phi_3|^2)$$

$\langle \Theta \rangle = \Theta_0$

$$\phi^\dagger \phi \Theta^\dagger X = \underbrace{\begin{pmatrix} \phi_1^\dagger \\ \phi_2^\dagger \\ \phi_3^\dagger \end{pmatrix}_3 \otimes \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}_3}_{2} \otimes \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}_2 \Theta^\dagger = \begin{pmatrix} \frac{1}{\sqrt{2}}(\phi_2^\dagger \phi_2 - \phi_3^\dagger \phi_3) \\ \frac{1}{\sqrt{6}}(-2\phi_1^\dagger \phi_1 + \phi_2^\dagger \phi_2 + \phi_3^\dagger \phi_3) \end{pmatrix}_2 \otimes \begin{pmatrix} X_1 \\ X_2 \end{pmatrix}_2 \Theta^\dagger = \frac{\Theta_0^* X_1}{\sqrt{2}} (|\phi_2|^2 - |\phi_3|^2)$$

$\langle X \rangle = (X_1, 0), \langle \Theta \rangle = \Theta_0$

## Vacuum structure

Potential

$$\begin{aligned}
 V &= -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 + c \phi^\dagger \phi X^\dagger X + k \phi^\dagger \phi \Theta^\dagger \Theta + (g \phi^\dagger \phi \Theta^\dagger X + h.c.) \\
 &= \frac{\Lambda_1}{2} (|\phi_1|^4 + |\phi_2|^4 + |\phi_3|^4) + \Lambda_2 (|\phi_1|^2 |\phi_2|^2 + |\phi_2|^2 |\phi_3|^2 + |\phi_3|^2 |\phi_1|^2) \\
 &\quad + \frac{\Lambda_3}{2} \left[ (\phi_1^\dagger \phi_2)^2 + (\phi_2^\dagger \phi_3)^2 + (\phi_3^\dagger \phi_1)^2 + h.c. \right] \\
 &\quad + \left( -\mu^2 + c_1 X_1^2 + \frac{2c_2 X_1^2}{\sqrt{6}} \right) |\phi_1|^2 + \left( -\mu^2 + c_1 X_1^2 + \frac{c_2 X_1^2}{\sqrt{6}} + \frac{g X_1 \Theta_0}{\sqrt{2}} \right) |\phi_2|^2 + \left( -\mu^2 + c_1 X_1^2 + \frac{c_2 X_1^2}{\sqrt{6}} - \frac{g X_1 \Theta_0}{\sqrt{2}} \right) |\phi_3|^2
 \end{aligned}$$

Potential minimum conditions

$$\left( \frac{\partial V}{\partial \phi_i} \right)_{\phi_1=v_1, \phi_2=v_2, \phi_3=v_3} = 0, \quad i = 1, 2, 3$$

vacuum expectation values

$$\begin{aligned}
 v_1 &= \pm \sqrt{\frac{\mu'^2}{\Lambda_1 + 2\Lambda'} - \frac{2\Lambda_1 c_2'}{(\Lambda_1 + 2\Lambda')(\Lambda_1 - \Lambda')}} \\
 v_2 &= \pm \sqrt{\frac{\mu'^2}{\Lambda_1 + 2\Lambda'} - \frac{\Lambda_1 - 2\Lambda'}{(\Lambda_1 + 2\Lambda')(\Lambda_1 - \Lambda')} c_2' - \frac{g'}{\Lambda_1 - \Lambda'}} \\
 v_3 &= \pm \sqrt{\frac{\mu'^2}{\Lambda_1 + 2\Lambda'} - \frac{\Lambda_1 - 2\Lambda'}{(\Lambda_1 + 2\Lambda')(\Lambda_1 - \Lambda')} c_2' + \frac{g'}{\Lambda_1 - \Lambda'}}
 \end{aligned}$$

Rewrite VEV with  
 $v$  and  $\alpha, \beta$

$$\langle \phi \rangle = \begin{pmatrix} v \sin \alpha \\ v \cos \alpha \sin \beta \\ v \cos \alpha \cos \beta \end{pmatrix}$$

$v$  : Higgs VEV  
 $\alpha, \beta$  : free parameter

## 7. Summary

We consider  $S_4$  symmetry as flavor symmetry.

We consider Higgs field  $\phi$  as  $S_4$  triplet.



We build new flavor model by using 3HDM and  $S_4$  symmetry.

We calculate mass matrices of charged leptons and neutrinos under new flavor model.

Mass matrix of charged leptons

$$M_l = M_{l1} + M_{l2}$$

$$= \begin{pmatrix} 0 & -\frac{2y_{e\mu}\theta_0}{\sqrt{6}\Lambda}v_1 & y_\tau v_1 \\ \frac{y_{e\mu}\theta_0}{\sqrt{2}\Lambda}v_2 & \frac{y_{e\mu}\theta_0}{\sqrt{6}\Lambda}v_2 & y_\tau v_2 \\ -\frac{y_{e\mu}\theta_0}{\sqrt{2}\Lambda}v_3 & \frac{y_{e\mu}\theta_0}{\sqrt{6}\Lambda}v_3 & y_\tau v_3 \end{pmatrix}_{LR} + \begin{pmatrix} \left(\frac{y_{l1}}{\Lambda} - \frac{2y_{l2}}{\sqrt{6}\Lambda}\right)v_1 X_1 & 0 & 0 \\ \left(\frac{y_{l1}}{\Lambda} + \frac{y_{l2}}{\sqrt{6}\Lambda}\right)v_2 X_1 & \frac{y_{l2}}{\sqrt{2}\Lambda}v_2 X_1 & 0 \\ \left(\frac{y_{l1}}{\Lambda} - \frac{y_{l2}}{\sqrt{6}\Lambda}\right)v_3 X_1 & -\frac{y_{l2}}{\sqrt{2}\Lambda}v_3 X_1 & 0 \end{pmatrix}_{LR}$$



We perform numerical analysis and calculate  $\delta_{CP}$ , effective mass  $m_{ee}$  and Majorana phases  $\eta_1, \eta_2$ .

We obtain strong predictions of  $\delta_{CP}$  and  $m_{ee}$  ( $m_{ee} \approx 0.0065[\text{eV}]$ ).

→ This flavor model can be confirmed by neutrino experiments in the near future.

Finally, we analysis the Higgs potential and we obtain the Higgs VEV  $\langle \phi \rangle = (v_1, v_2, v_3)$ .

Mass matrix of left-handed Majorana neutrinos

$$m_\nu = -M_D M_R^{-1} M_D^\dagger$$

$$M_D = \begin{pmatrix} y_{De}v_1 & 0 & -2/\sqrt{6}y_{D\mu\tau}v_1 \\ y_{De}v_2 & 1/\sqrt{2}y_{D\mu\tau}v_2 & 1/\sqrt{6}y_{D\mu\tau}v_2 \\ y_{De}v_3 & -1/\sqrt{2}y_{D\mu\tau}v_3 & 1/\sqrt{6}y_{D\mu\tau}v_3 \end{pmatrix}_{LR} \quad M_R = \begin{pmatrix} M_{eR} & 0 & 0 \\ 0 & M_{\mu\tau R} & 0 \\ 0 & 0 & M_{\mu\tau R} \end{pmatrix}$$