

## Pulsed laser spectroscopy of Muonium 1S-2S transition in J-PARC (#6)

*Thursday, 14 March 2024 13:20 (10 minutes)*

“Our purpose is precision measurement of the 1S-2S energy interval in Muonium, which is an exotic hydrogen-like atom consists of a positive muon and an electron. This purely leptonic system enables a precise calculation of the energy interval with the Standard Model without any concerns of the uncertainty from the charge radius of the nucleus, unlike the hydrogen atom. This advantage motivates us to measure the precise 1S-2S energy interval in Muonium with technology of laser spectroscopy and to determine the muon mass with the highest accuracy of 10 ppb. The improvement of muon mass accuracy has an impact on verification of the Standard Model, muon  $g-2$ /EDM experiment, for example.

We will report a recent result of Muonium 1S-2S energy interval measurement at J-PARC. The event rate of 1S $\rightarrow$ 2S in our experiment is 50 times higher than the previous experiment[1]. This improvement gives promising prospect for higher accuracy of 1S-2S energy interval and the muon mass accuracy in the future.

1, Meyer, V. et al. Measurement of the 1s-2s energy interval in muonium. Phys. Rev. Lett. 84, 1136 (2000).”

**Presenter:** YAMAMOTO, Shinsuke (Okayama University)

**Session Classification:** Poster