

Quantum computing of chirality imbalance in $SU(2)$ gauge theory

Xingyu Guo (QuNu Collaboration)
South China Normal University

arXiv:XXXX.XXXXX

Introduction

- Spontaneous chiral symmetry breaking
 - One of the key features of QCD
 - Origin of mass
 - Chiral magnetic effect, chiral vortical effect...
 - Non-perturbative, high baryon chemical potential
 - Challenging for traditional methods

Introduction

- Quantum computing: a promising new method
- Topics of interest:
 - Real-time evolution
 - Non-perturbative physics
 - Non-Abelian gauge theory
 - Thermal states
- ...

Model

- 1+1D SU(2) model: simplest non-Abelian model

$$H = -i\bar{\psi}\gamma^1(\partial_1 + igA_1^a t^a)\psi + m\bar{\psi}\psi + \mu\psi^\dagger\psi + \frac{1}{2}\sum_a (L^a)^2$$

- Chiral condensate: $\sigma = \bar{\psi}\psi$
- Discretization: Staggered fermion

$$\psi_1(x) \rightarrow \phi_{2n}, \quad \psi_2(x) \rightarrow \phi_{2n+1}$$

$$H = \frac{1}{2\Delta} \sum_{n=0}^{N-2} (\phi_n^\dagger U_n \phi_{n+1} + H.C.) + m \sum_{n=0}^{N-1} (-1)^{n+1} \phi_n^\dagger \phi_n + \mu \sum_{n=0}^{N-1} \phi_n^\dagger \phi_n + \frac{\Delta g^2}{2} \sum_{n=0}^{N-2} \mathbf{L}_n^2$$

Eliminating Gauge Field

- Gauss's Law

$$L_n^a - R_{n-1}^a = Q_{n-1}^a = \phi_n^\dagger t^a \phi_n \rightarrow L_n^a = \sum_{i < n} Q_i^a$$

- Local gauge transformation

$$\Theta = \prod_k \exp \left(i\theta_k \cdot \sum_{j > k} \mathbf{Q}_j \right)$$

$$H = \frac{1}{2} \sum_{n=0}^{N-2} (\phi_n^\dagger \phi_{n+1} + \text{H.C.}) + \Delta m \sum_{n=0}^{N-1} (-1)^{n+1} \phi^\dagger \phi_n + \Delta \mu \sum_{n=0}^{N-1} \phi^\dagger \phi_n + \frac{\Delta^2 g^2}{2} \sum_{n=0}^{N-2} \left(\sum_{k \leq n} \mathbf{Q}_k \right)$$

Jordan-Wigner transformation

$$\phi_n^r \rightarrow \varphi_{2n}, \quad \phi_n^g \rightarrow \varphi_{2n+1}$$

$$\varphi_n = \sigma_n^- \prod_{l=0}^{n-1} (-i\sigma_l^z)$$

$$H = -\frac{1}{2} \sum_{n=0}^{N-2} \left(\sigma_{2n}^+ \sigma_{2n+1}^z \sigma_{2n+2}^- + \sigma_{2n+1}^+ \sigma_{2n+2}^z \sigma_{2n+3}^- + \text{H.C.} \right) + \Delta m \sum_{n=0}^{N-1} \left[(-1)^{n+1} \frac{\sigma_{2n}^z + \sigma_{2n+1}^z}{2} + 1 \right] + \frac{\Delta^2 g^2}{2} \sum_{n=0}^{N-2} \left(\sum_{k \leq n} \mathbf{Q}_k \right)$$

$$Q_n^x = \frac{1}{2} (\sigma_{2n+1}^+ \sigma_{2n}^- + \text{H.C.})$$

$$Q_n^y = \frac{i}{2} (\sigma_{2n+1}^+ \sigma_{2n}^- - \text{H.C.})$$

$$Q_n^z = \frac{1}{4} (\sigma_{2n}^z - \sigma_{2n+1}^z)$$

Algorithm: Variational Method

- Finite temperature: the Gibbs state

$$\rho(\beta) = \frac{1}{Z(\beta)} e^{-\beta H}, \quad Z(\beta) = \text{Tr}(e^{-\beta H})$$

$$\rho(\alpha) = \sum_i P_i(\beta) |\varphi_i\rangle\langle\varphi_i|$$

Analytically solvable


$$P_i = \frac{e^{-\beta E_i}}{\left(\sum_n e^{-\beta E_n}\right)}$$

Eigenstate of H
independent of β

- Variational method
 - Parametrization

$$\rho(\alpha) = \sum_i P_i(\beta) U(\alpha) |i\rangle\langle i| U^\dagger(\alpha)$$

Algorithm Workflow



Algorithm: Variational Method

- Loss Function : Free energy

$$\begin{aligned} F &= E - TS \\ &= \text{Tr} [\rho H] + T \text{Tr} [\rho \log \rho] \\ &= \sum_i P_i [E_i + T \ln P_i] \end{aligned}$$

- The variational method is only used once for all different temperatures.
- Many part of the calculation is analytical.

Algorithm: Variational Method

- Construction of $U(\alpha)$: QAOA ansatz

$$U(\alpha) = \prod_{i=1}^p \prod_{j=1}^n \exp(i\alpha_{ij} H_j)$$

$$H = H_1 + H_2 + \cdots + H_n$$

$$[H_i, H_j] \neq 0$$

- Each H_i preserves the same symmetries as H , so that $U(\alpha)$ also preserves the symmetries.

Algorithm: Monte-Carlo


- Monte-Carlo in optimization
 - Randomly select a small set of states, do the optimization with this set until finished.
 - Select another set, continue the optimization with the new set.
 - Repeat until parameters convergence.
- In practical, a set of 20 qubits is used for each step.

Algorithm: Monte-Carlo

- Monte-Carlo in thermal state construction
 - Start from $|i\rangle$ such that $U|i\rangle$ is the ground state.
 - Randomly flip one qubit of $|i\rangle$ to get a new state $|j\rangle$
 - Calculate the energy expectation value $E_j \langle i | U^\dagger H U | i \rangle$
 - If $E_j < E_i$, accept the new state, otherwise, accept it with the probability $e^{-(E_j - E_i)/T}$
 - If the new state is rejected, the old state is added into the mixed state again.
 - Repeat until number of states reaches a predetermined limit.

Results: Full Gibbs State


- 8 qubits, $\mu = 0$.
- Optimization done at highest temperature.
- All 256 states are used to construct the Gibbs state.
- The VQE method produces the Gibbs state very accurately.



Chiral condensate at finite temperature

Results: Full Gibbs State


- 8 qubits, $m/g = 5$.
- The ground state changes at high chemical potential.
- Consistent with theoretical prediction.



Chiral condensate at finite temperature

Results: Monte-Carlo


- 8 qubits, 1000 states for each sampling.
- The accuracy is good.
- Monte-Carlo method is not effective for small system. (1000 vs. 256)



Chiral condensate at finite temperature

Results: Monte-Carlo


- 12 qubits, 1000 (left) and 2000 (right) states.
- Required number of sampling increases only as power law of number of qubits.
- 2000 vs. 4096 is already effective.



Chiral condensate at finite temperature

Results: Real QC results


- 8 qubits, results from IBM's quantum hardware.
- A simpler ansatz for U is used.
- Optimization is still done classically.
- No error mitigation used.
- Our algorithm can achieve good precision on real QC.



Chiral condensate at finite temperature

Results: Real QC results

- Good accuracy for all the eigenenergy.
- Promising to apply to larger systems.



Relative error of the eigenenergy

Summary and Outlook

- We propose a framework with VQE and Monte-Carlo method to simulate thermal states on quantum computers.
- With this frame work, the chiral condensate of 1+1D SU(2) gauge model is studied.
- Our method is efficient and accurate in classical simulations as well as on real QCs.
- Apply to massless fermions: spontaneous symmetry breaking. (Paper to appear soon)
- Extend to larger systems or even higher dimensions.