

B physics: known and unknown

Zoltan Ligeti

KEK Theory Meeting on Particle Physics Phenomenology

February 16 – 19, 2026



Outline

- Why flavor?
(Unique sensitivities)
- Recent examples of what we know and don't know
(Lot to do, both for theorists and experimentalists)
- Future: HL-LHC & Belle II sensitivity \Rightarrow then FCC
(Huge increases in data, what can they teach us?)

What is flavor physics?


- Flavor \equiv what distinguishes the three generations? (break $[U(3)]^5$ global sym.)
Experimentally: rich and sensitive ways to probe the SM, and search for NP
- **SM flavor**: masses? mixing angles? 3 generations? — most of the SM parameters
Flavor in SM is simple: only from Yukawa couplings to Higgs, **many testable relations**
- **BSM flavor**: TeV scale (hierarchy puzzle) \ll “naive” scale of flavor & CP violation
New particles that couple to quarks or leptons \Rightarrow new flavor param's, **clues about BSM**
- **Baryon asymmetry** requires CPV beyond the SM — how precisely can we test it?
(Not necessarily in flavor changing processes, nor necessarily in the quark sector)
- **Neutrino mass**: What is the Lagrangian? (is lepton # violated?)

Origin of high scale sensitivity


- Large suppressions in the SM — back of an envelope calculation of Δm_K :
- Why is $\Delta m_K/m_K \sim 7 \times 10^{-15}$?

In the SM:
$$\frac{\Delta m_K}{m_K} \sim \alpha_w^2 |V_{cs}V_{cd}|^2 \frac{m_c^2}{m_W^4} f_K^2$$

Predicted $m_c \sim 1.5 \text{ GeV}$ (Gaillard & Lee; Vainshtein & Khriplovich, 1974)



- If exchange of a heavy particle X contributes at the SM level:



$$\frac{\Delta m_K^{(X)}}{\Delta m_K} \sim \frac{g^2 \Lambda_{\text{QCD}}^3}{M_X^2 \Delta m_K} \Rightarrow M_X > g \times 10^3 \text{ TeV}$$

- Alternatively, sensitivity to TeV-scale particles with one-loop couplings [$g \sim \mathcal{O}(10^{-3})$]

Similar story for many other flavor-changing neutral current (FCNC) processes

New physics scales probed

- Heavy new physics generates dimension > 4 operators in SMEFT

$$\mathcal{L} = \text{SM} + \sum_i \frac{C_{5i}}{\Lambda} \mathcal{O}_{5i} + \sum_i \frac{C_{6i}}{\Lambda^2} \mathcal{O}_{6i} + \dots$$

- Dim-5 $LL\phi\phi$ discovered, iff m_ν violates lepton # ($0\nu\beta\beta?$); No dim-6 term established yet

- Baryon number violation: $\frac{QQQL}{\Lambda^2} \Rightarrow \Lambda \gtrsim 10^{16} \text{ GeV}$

- Higgs & precision electroweak: $\frac{(\phi D^\mu \phi)^2}{\Lambda^2} \Rightarrow \Lambda \gtrsim 10^4 \text{ GeV}$

- Flavor and CP violation (and EDMs): $\frac{QQQQ}{\Lambda^2} \Rightarrow \Lambda \gtrsim 10^{(3\dots 8)} \text{ GeV}$

The only experimental probe of the $10^2 - 10^5 \text{ TeV}$ scale

Compare flavor and precision electroweak

- Flavor-changing neutral currents

No tree level

Loop + CKM + GIM + CPV suppression

$$\frac{1}{\Lambda^2} \sim \frac{\alpha_W}{m_W^2} (V_{ik} V_{jk}^*) \frac{m_q^2}{m_W^2} \frac{J_{\text{CKM}}}{|V_{ik} V_{jk}|}$$

Probe $\Lambda \sim 10^5 \text{ TeV}$

- EW precision measurements

Tree level



Loop suppression

$$\frac{1}{\Lambda^2} \sim \frac{\alpha_W}{m_W^2}$$

Probe $\Lambda \sim 10 \text{ TeV}$

- Both are important: new physics probably has additional small parameters / structures



Plots from European strategy



Some lessons: not new, but got lot sharper


- Flavor was critical in developing the SM, and mostly an input to model building
- All TeV-scale NP models must contain some mechanism to avoid violating constraints
E.g., in SUSY, to suppress flavor and CP violation, impose: universality, heavy squarks, alignment
Devised to make deviations small from the SM, revisit model building when tensions with SM arise
- The SM suppressions are strongest for kaons (1st – 2nd generation)
For many models, Δm_K and ϵ_K are the most constraining — 3rd gen. “looks” different
The idea of (dominantly) 3rd generation NP goes back (at least) to the mid-90s [[hep-ph/9512388](#), [hep-ph/9607394](#)]
- What can future data tell us?
If deviations from SM, upper bound on BSM scale, help understand its structure

Large upcoming improvements



- LHCb upgrade in LS2 (inst. lumi.: 2×10^{33})
- LHCb Upgrade II in LS4 (inst. lumi.: 1.5×10^{34})
- ATLAS & CMS competitive in some modes
- Belle II goal: over $50 \times$ Belle data set
- Discussions about physics & feasibility of an upgrade (polarization? 250/ab?)

Extensive sensitivity projections: LHCb Upgrade II [1808.08865]; Belle II [1808.10567]; and including ATLAS & CMS @ HL-LHC [2503.24346]



Many uncertainties will improve by 5–10

Where are we now? Testing quark flavor


- $(u, c, t) W^\pm (d, s, b)$: 9 complex couplings \Rightarrow many relations Are they consistent?

$$V_{\text{CKM}} = \underbrace{\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}}_{\text{CKM matrix}} = \begin{pmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + \dots$$

Measurements: magnitudes \sim decay rates; phases $\sim CP$ violation


- Many observables are $f(\rho, \eta)$ — want to compare:

- $b \rightarrow ul\bar{\nu} \Rightarrow |V_{ub}/V_{cb}|^2 \propto \rho^2 + \eta^2$
- $\Delta m_{B_d}/\Delta m_{B_s} \Rightarrow |V_{td}/V_{ts}|^2 \propto (1 - \rho)^2 + \eta^2$
- CP violation in K, B, B_s decays




The CKM fit in the SM

- Spectacular progress in last 25 years
- The CKM mechanism describes consistently CP violation and flavor changing processes



The CKM fit in the SM

- Spectacular progress in last 25 years
 - The CKM mechanism describes consistently CP violation and flavor changing processes
 - Looser constraints if NP is allowed in fits
- Full fit (upper plot) vs. tree-level (lower plot)
- LHCb: B_s constraints caught up with B_d
 - $\mathcal{O}(20\%)$ NP contributions to most loop-level processes (FCNC) are still allowed




Constraints on NP parameters added to the SM


- In a large class of NP models, dominant contributions are to neutral meson mixing

- General parametrization of many scenarios by two real BSM parameters; redo CKM fit:

$$h e^{2i\sigma} = A_{\text{NP}}(B^0 \rightarrow \bar{B}^0) / A_{\text{SM}}(B^0 \rightarrow \bar{B}^0)$$



$$\text{SM: } \frac{C_{\text{SM}}}{m_W^2}$$






$$\text{NP: } \frac{C_{\text{NP}}}{\Lambda^2}$$

Tree-level decays dominated by SM, BSM only significant in FCNCs (loops)

- Is $h \gtrsim 1$ allowed? If not, then CKM mechanism is dominant

Evolving bounds on $h_{d,s}$

- CKM fit with $h_{d,s}$ and $\sigma_{d,s}$ parameters added: $h e^{2i\sigma} = A_{\text{NP}}(B^0 \rightarrow \bar{B}^0) / A_{\text{SM}}(B^0 \rightarrow \bar{B}^0)$



[color: 2σ , dotted: 3σ]

- LHCb: BSM contributions to B_s mixing are even more constrained than those to B_d
- Weak interaction dominates observed CP violation: $\text{BSM} / \text{SM} \lesssim 20\%$


Improvements in next ~ 10 years

- At 95% CL: $NP \lesssim (0.25 \times SM) \Rightarrow NP \lesssim (0.08 \times SM)$

- Scale: $h \simeq \frac{|C_{ij}|^2}{|V_{ti}^* V_{tj}|^2} \left(\frac{4.5 \text{ TeV}}{\Lambda} \right)^2$

$$\Rightarrow \Lambda \sim \begin{cases} 2.3 \times 10^3 \text{ TeV} \\ 20 \text{ TeV (tree + CKM)} \\ 2 \text{ TeV (loop + CKM)} \end{cases}$$

- Complementary to high- p_T searches (E.g., similar to HL-LHC $m_{\tilde{g}}$ reach)
- BSM sensitivity will continue to improve



[color: 2σ , dotted: 3σ]

$$h e^{2i\sigma} = A_{NP}(B^0 \rightarrow \bar{B}^0) / A_{SM}(B^0 \rightarrow \bar{B}^0)$$

Recent tensions between SM and data

Hints of deviations from the SM

- From a distance, everything seems to work... looking closer:
 - Charged current semileptonic decays (e.g., $B \rightarrow D^{(*)} \ell \bar{\nu}$)
 - Neutral current (FCNC) semileptonic decays (e.g., $B \rightarrow K^{(*)} \ell \bar{\ell}$)
 - Dominantly charged current nonleptonic decays (e.g., $B \rightarrow D^{(*)} h$)
 - Dominantly FCNC nonleptonic decays (e.g., $B \rightarrow K h$)

There are more...

- Intriguing tensions with the SM \Rightarrow experimental scrutiny, new theory ideas
- What are smallest deviations from SM, which can be unambiguously established?

The $B \rightarrow D^{(*)}\tau\bar{\nu}$ decay rates

- BaBar, Belle, LHCb: $R(X) = \frac{\Gamma(B \rightarrow X\tau\bar{\nu})}{\Gamma(B \rightarrow X(e/\mu)\bar{\nu})}$


$\sim 3.5\sigma$ from SM expectations: theory robust due to heavy quark symmetry + lattice QCD

- Imply NP scale \lesssim few TeV



Mediators within / near ATLAS & CMS reach

- Tree level: three ways to insert mediator: $(b\nu)(c\tau)$, $(b\tau)(c\nu)$, $(bc)(\tau\nu)$
overlap with ATLAS & CMS searches for \tilde{b} , leptoquark, H^\pm

- Models built to fit these impacted ATLAS & CMS searches, motivated LFV searches



Large future improvements



[2101.08326]

- Measurements will improve a lot, and reach few % in several decay modes
- May establish NP, even if deviations from SM decrease
- Will at least lead to much more robust $|V_{cb}|$ — critical for precision in many observables

$|V_{cb}|$ and pushing HQET further

- $|V_{cb}|$ is measured from “zero recoil” limit of $d\Gamma(B \rightarrow D^{(*)} \ell \bar{\nu})/dq^2$

HQET most constraining + LQCD most precise

In practice, measuring all distributions helps — additional information

Impacts many rare decay rate predictions, since $V_{ts}, V_{td} \propto A$


- Many open questions:

How to fit data: truncate model independent (BGL) param.? (Many proposals, Akaike information criterion, etc.)

Optimal combination with constraints from unitarity and lattice QCD?

Some tension between LQCD and data



- Not to mention tension between exclusive and inclusive determination



[Bernlochner, ZL, Persson, Robinson, to appear]


R_K and R_{K^*} until 2022

- LHCb: $R_{K^{(*)}} = \frac{B \rightarrow K^{(*)} \mu^+ \mu^-}{B \rightarrow K^{(*)} e^+ e^-} < 1$ three bins $\sim 2.5\sigma$ from lepton universality







- Theorists' fits quoted $3-5\sigma$ (sometimes including P'_5 and/or $B_s \rightarrow \phi \mu^+ \mu^-$)
- Modifying one Wilson coefficient in \mathcal{H}_{eff} gave good fit: $\delta C_{9,\mu} \sim -1$

R_K and R_{K^*} now: SM-like, but rates too small?



$0.1 < q^2 < 1.1 \text{ GeV}^2$ (low- q^2), $1.1 < q^2 < 6.0 \text{ GeV}^2$ (central)



[Smith, LHCP 2024]

The P_5' angular distribution in $B \rightarrow K^* \mu^+ \mu^-$

- “Optimized observables” [1202.4266 + long history]
(assumptions about factorizable / nonfactorizable)


Global fits: best fit, still: NP reduces C_9

[Altmannshofer, Straub; Descotes-Genon, Matias, Virto; Jager, Martin Camalich; Bobet, Hiller, van Dyk; many more]

- Difficult for lattice QCD, large recoil

What calculation determines how far below $m_{J/\psi}$ this comparison is reliable? (Different than $e^+e^- \rightarrow$ hadrons)


- Tests: other observables, q^2 dep., B_s and Λ_b decays
- Is the $c\bar{c}$ loop tractable? Impacts many interesting decay rates & CP viol. observables



BSM, fluctuation, SM theory?


Intense discussions of theory uncertainties

- My hope: data will help to make progress
Large data sets often inspired progress in theory, and developing new approaches
- Only robust deviations based on model-independent theory will be seen as signs of new physics



$B \rightarrow K \nu \bar{\nu}$ — unique to e^+e^- colliders


- Similar short-distance contributions, and much simpler long-distance ones



- Also relevant for dark sector searches ($B \rightarrow K + \text{invis.}$)
(Is the excess in one bin of q^2 ?) [2309.00075, 2311.14629, 2602.09666]

- Input: precise form factor calculation [HPQCD, 2207.12468]


- If this tension becomes more significant, stopping NA62 after LHC Run 3 will look even more mistaken



Belle II: 2.7σ from SM

Nonleptonic $b \rightarrow c$ decays

- Some $B \rightarrow Dh$ decays factorize in heavy quark limit + power suppressed corrections



[LHCb, 2/10/2026: direct CPV in $B_s^0 \rightarrow D_s^- \pi^+ \lesssim 1\%$]

- Need: semileptonic form factor near $q^2 = 0$, estimate power suppressed corrections

(Still) not understood: the $B \rightarrow K\pi$ puzzle

- Have we seen new physics in CPV?

$$A_{K^+\pi^-} = -0.0831 \pm 0.0031 \quad (P + T)$$


$$A_{K^+\pi^0} = 0.027 \pm 0.012 \quad (P + T + C + A + P_{ew})$$

- Large difference — small SM sources?

$$A_{K^+\pi^0} - A_{K^+\pi^-} = 0.111 \pm 0.012$$

SCET / factorization predicts: $\arg(C/T) = \mathcal{O}(\Lambda_{\text{QCD}}/m_b)$ and $A + P_{ew}$ small

- Large fluctuations? Breakdown of $1/m$ expansion? Missing something subtle? BSM?
- Can we unambiguously understand theory, so that such data could disprove the SM?



Belle II can make unique contribution

- Sum rule based on isospin for dominant amplitudes, and $SU(3)$ for subleading terms:


$$\delta_{K\pi} = \Delta(K^+\pi^-) + \Delta(K^0\pi^-) - 2\Delta(K^+\pi^0) - 2\Delta(K^0\pi^0) \quad \Delta(f) = \Gamma(\bar{B} \rightarrow \bar{f}) - \Gamma(B \rightarrow f)$$

Expect: $|\delta_{K\pi}/\Delta(K^+\pi^-)| \lesssim \text{few } \%$

[Gronau, hep-ph/0508047]

$$I_{K\pi} = \frac{\delta_{K\pi}}{2\Gamma(B \rightarrow K^+\pi^-)}$$

Interesting to measure well beyond
Belle II sensitivity \Rightarrow FCC- ee



(Far) future


$\sin 2\beta$ and $B_q \rightarrow J/\psi h$

$(q = u, d, s; \quad h = \pi, K)$

Details: ZL, Y. Nir, R. Schein, JHEP 10 (2024) 100, 2506.21675

Towards theory at the per mille level

- Theory uncertainty of γ is negligible (2nd order EW)
World average $\gamma = (66.4 \pm 2.8)^\circ$
- Theory uncertainty of “everything else” being discussed
- Most likely, β will be second best
There are claims of possibly large $|V_{ub}|$ (“penguin”) contamination
- World average: $\sin 2\beta = 0.710 \pm 0.011 \quad (22.62 \pm 0.45)^\circ$



16 observables in $B_q \rightarrow J/\psi h$

Process	\mathcal{B}	A or C	S
$B^+ \rightarrow J/\psi \pi^+$	$(3.92 \pm 0.09) \times 10^{-5}$	$(+1.23 \pm 0.47) \times 10^{-2}$	—
$B^+ \rightarrow J/\psi K^+$	$(1.020 \pm 0.019) \times 10^{-3}$	$(+1.8 \pm 3.0) \times 10^{-3}$	—
$B_d \rightarrow J/\psi \pi^0$	$(1.74 \pm 0.07) \times 10^{-5}$	$(+8.5 \pm 8.5) \times 10^{-2}$	-0.87 ± 0.11
$B_d \rightarrow J/\psi K^0$	$(8.91 \pm 0.21) \times 10^{-4}$	$(+0.9 \pm 1.0) \times 10^{-2}$	$+0.708 \pm 0.012$
$B_s \rightarrow J/\psi \pi^0$	$< 1.21 \times 10^{-5}$	NA	NA
$B_s \rightarrow J/\psi K_S$	$(1.92 \pm 0.14) \times 10^{-5}$	$-0.28 \pm 0.41 \pm 0.08$	$-0.08 \pm 0.40 \pm 0.08$

Recent experimental results: LHCb, 2411.12178, 2402.05528; Belle II 2410.08622; Belle 2311.12724

12 parameters in $B_q \rightarrow J/\psi h$

● Expand amplitudes: $A = (V_{cb}^* V_{cq}) (A_c^{(0)} + \varepsilon A_c^{(1)}) + (V_{ub}^* V_{uq}) A_u^{(0)} + \dots$

● $SU(3)$ flavor symmetry \Rightarrow 5 $A_c^{(0)}$, 4 $A_c^{(1)}$, 3 $A_u^{(0)}$ relations

[ZL & Robinson, 1507.06671]

● 6 independent complex amplitudes: 1 $A_c^{(0)}$, 2 $A_c^{(1)}$, 3 $A_u^{(0)}$

So there must be 4 relations when working to this order

4 relations for $B_q \rightarrow J/\psi h$ at this order

- $A_{\psi K^+}/A_{\psi \pi^+} = -|V_{cd}/V_{cs}|^2$
 \Rightarrow LHCb's ΔA_{CP} implies: $A_{\psi \pi^+} \simeq (1.23 \pm 0.47) \times 10^{-2}$ and $A_{\psi K^+} \simeq (-6.5 \pm 2.5) \times 10^{-4}$
- $C_{\psi K^0}^d/C_{\psi \bar{K}^0}^s = -|V_{cd}/V_{cs}|^2$
 \Rightarrow Narrow allowed range of $C_{\psi \bar{K}^0}^s = -0.17 \pm 0.19$ compared to direct measurement
- $(1 + \bar{\lambda}^2)s_{2\beta} = S_{\psi K_S}^d - \bar{\lambda}^2 S_{\psi \pi}^d - 2(\Delta_K + \bar{\lambda}^2 \Delta_\pi)c_{2\beta} \tan \gamma + \mathcal{O}(\varepsilon R_u \bar{\lambda}^2, R_u^2 \bar{\lambda}^2, \delta)$ [1507.06671]
where: $\Delta_K = \frac{\Gamma(B_d \rightarrow \psi K^0) - \Gamma(B^+ \rightarrow \psi K^+)}{\Gamma(B_d \rightarrow \psi K^0) + \Gamma(B^+ \rightarrow \psi K^+)}$, $\Delta_\pi = \frac{2\Gamma(B_d \rightarrow \psi \pi^0) - \Gamma(B^+ \rightarrow \psi \pi^+)}{2\Gamma(B_d \rightarrow \psi \pi^0) + \Gamma(B^+ \rightarrow \psi \pi^+)}$
 $\Rightarrow \sin 2\beta - S_{\psi K_S}^d = 0.05 \pm 0.03$ (+ issues of production asym vs. isospin violation [2306.04686])
- $S_{\psi K_S}^d - s_{2\beta} = -|V_{cd}/V_{cs}|^2 (c_{2\beta}/c_{2\beta_s}) (S_{\psi K_S}^s + s_{2\beta_s})$ [2506.21675]
 $\Rightarrow S_{\psi K_S}^d - \sin 2\beta = 0.001 \pm 0.015$

Extracting $\sin 2\beta$

- Substantial range of predictions in literature for $S_{\psi K_S}^d - \sin 2\beta$
- Neglecting $|(V_{ub}V_{us})/(V_{tb}V_{ts})(P/T)|$ uncertainty should be $\mathcal{O}(0.01)$

$$\sin 2\beta = S_{\psi K_S}^d$$

- Using flavor $SU(3)$, uncertainty should be $\mathcal{O}(0.003)$



$$\sin 2\beta = S_{\psi K_S}^d + \left| \frac{V_{cd}}{V_{cs}} \right|^2 \frac{c_{2\beta}}{c_{2\beta_s}} (S_{\psi K_S}^s + s_{2\beta_s})$$

- At next order: hadronic uncertainty suppressed if $C_{\psi K_S}^s$ is small

$$\sin 2\beta = S_{\psi K_S}^d + \left| \frac{V_{cd}}{V_{cs}} \right|^2 \frac{c_{2\beta}}{c_{2\beta_s}} (S_{\psi K_S}^s + s_{2\beta_s}) \left[1 + \frac{R_{K^0 K^0}^{sd} - 1}{2} + \frac{C_{\psi K_S}^s c_{2\beta_s}}{S_{\psi K_S}^s} \left(\varepsilon \operatorname{Im} \frac{A_c^{(1)}}{A_c^{(0)}} - \rho(1 + \bar{\lambda}^2) \frac{C_{\psi K_S}^s}{2\eta} \right) \right]$$

- **Wishlist: More precise $S_{\psi K_S}^s$ and $C_{\psi K_S}^s$** (LHCb result only used 3/fb in Run 1 [1503.07055])

Flavor @ FCC



● Of the future colliders discussed, only FCC would go well beyond LHCb + Belle II goals

Flavor physics at FCC- ee

- $10^5 \times \text{LEP}$ is the right target (mass scale) \propto (uncertainty) $^{-1/2} \propto$ (stat) $^{-1/4}$
- Can one appreciate / anticipate a 10^5 improvement? (Recall: Belle II / ARGUS $\sim 10^5$)
Theory and experimental techniques both changed a lot! (e.g., full hadronic reconstruction)
Asymmetric B factories at $\Upsilon(4S)$ great for CP violation, less ideal for (semi)leptonic decays
- What was not even tried at LEP? (due to lack of statistics or lack of physics interest)
- Some rare decay sensitivity may improve linearly with statistics; e.g., $Z \rightarrow \mu\tau, \mu e$, etc.
- A lot of what's usually called precision electroweak, also concern flavor (τ lifetime & mass, R_b, R_ℓ for each ℓ flavor, etc.)

FCC- ee data sets for flavor

- Production yields compared to Belle II: [2106.01259]

Particle production (10^9)	$B^0 + \bar{B}^0$	B^\pm	$B_s^0 + \bar{B}_s^0$	$\Lambda_b + \bar{\Lambda}_b$	B_c^\pm	$c\bar{c}$	$\tau^+\tau^-$
Belle II (50 ab^{-1})	27	27	tbd	—	—	65	45
tera- Z ($5 \times 10^{12} Z$)	600	600	150	130	3	600	170

Comparison with LHCb more complex: roles of trigger, LHCb has advantage if final state is fully reconstructed, if there are neutrals, tera- Z may win

- WW threshold ($10^8 WW$): $W \rightarrow b\bar{c}$ can give a qualitatively new determination of $|V_{cb}|$
Estimate 0.2% uncertainty, independent of B decays [Monteil @ 7th FCC Workshop, Jan 2024]; [2405.08880]
- Some electroweak precision measurements also concern flavor (R_ℓ for each flavor, τ_τ, m_τ)

Sensitivity to new physics in B mixing



- In many BSM scenarios, dominant deviations from SM may be in neutral meson mixing

Assume: (i) 3×3 CKM matrix is unitary; (ii) tree-level decays dominated by SM

General parametrization: $h e^{2i\sigma} = A_{\text{NP}}(B^0 \rightarrow \bar{B}^0)/A_{\text{SM}}(B^0 \rightarrow \bar{B}^0)$ ($h_{d,s}, \sigma_{d,s}$: NP param's)


- CKM fit with 4 BSM param's added; combines many measurements and theory inputs [Charles *et al.*, 2006.04824] (\Rightarrow conservative view of future progress)

- $|V_{cb}|$ becomes a bottleneck; Tera- Z sensitivity will be better (no LQCD extrapolations)



If BSM in $b \rightarrow c\tau\bar{\nu}$, must study $\nu\bar{\nu}$ & $\tau\tau$ modes



- SM and data in 3σ tension in $R(D^{(*)})$, if established, requires $\mathcal{O}(10\%)$ correction to a tree-level SM process
- If NP is charged under $SU(2)$, **unavoidable connection** to $b \rightarrow s\tau^+\tau^-$ or $b \rightarrow s\nu\bar{\nu}$ — correlations distinguish models



Only Z can measure $B \rightarrow K^*\tau^+\tau^-$, $K^*\nu\bar{\nu}$ at SM level

Belle II: $\mathcal{B}(B \rightarrow K^*\tau^+\tau^-) < 1.8 \times 10^{-3}$ (SM $\sim 10^{-6}$) [2504.10042]

- Boost of B in Z decay provides ideal environment



$B_{d,s} \rightarrow \mu^+ \mu^-$: interesting at all imaginable statistics

- $\mathcal{B}_{\text{SM}}(B_s \rightarrow \mu^+ \mu^-) \sim 3 \times 10^{-9}$, BSM predictions extended orders of magnitudes higher
- $\mathcal{B}_{\text{SM}}(B_d \rightarrow \mu^+ \mu^-) \sim 10^{-10}$, LHCb expects 10% (300/fb), CMS expects 15% (3/ab)


SM uncertainty $\simeq (2\%) \oplus f_{B_q}^2 \oplus \text{CKM}$, and may be further reduced

- Measure $|V_{ub}|$, using only isospin, from:

$$\mathcal{B}(B_u \rightarrow \ell \bar{\nu}) / \mathcal{B}(B_d \rightarrow \mu^+ \mu^-)$$

(Most precise $|V_{ub}|$... at FCC- hh statistics...?)

- A decay with mass-scale sensitivity (dim.-6 operator) that competes with $K \rightarrow \pi \nu \bar{\nu}$




PDG average:

$$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-) = (3.34 \pm 0.27) \times 10^{-9}$$

Actual discovery potential? E.g., SUSY in $Z \rightarrow \ell^+ \ell^-$


- Consider a SUSY simplified model: \tilde{q}, \tilde{g} heavy, only electroweakinos & sleptons light

- Measurement: $R_\ell = \frac{\Gamma_{\ell^+ \ell^-}}{\Gamma_{\text{hadrons}}}$
- | | | | | |
|--------------------------|---|-------------|------------------------|-----------------------------|
| $R_\ell^Z (\times 10^3)$ | 20767 ± 25 | 0.05 | 0.05 | Ratio of hadrons to leptons |
| | Systematic limitation in Midterm Report resolved by Feasibility Study | | Acceptance for leptons | |



- Ultimate sensitivity depends on $\alpha_s, \sin^2 \theta_w$, etc. Several measurements combined for best physics reach Even better sensitivity to flavor violating effects ($e/\mu/\tau$)

- Can probe beyond (or between) HL-LHC exclusions
- Complementary to SMEFT studies, a specific model may have important correlations



Final remarks

What are the largest useful data sets?

- No one has seriously explored it!
- Many measurements will remain far from being limited by theory uncertainties:
 - For $\gamma \equiv \phi_3$, theory uncertainty only from higher order EW
 - $B_{s,d} \rightarrow \mu\mu$, $B \rightarrow \mu\nu$ and other leptonic decays (lattice QCD, [double] ratios)
 - $A_{\text{SL}}^{d,s}$ — will experimental systematics become limiting?
 - Lepton flavor violation & lepton universality violation searches
 - Probably many more...
- Very broad program, dark sector searches, etc.
- In some decays, even in 2040s we'll have (exp. bound)/SM $\gtrsim 10^3$ (E.g., $B_{d,s} \rightarrow e^+e^-$, $\tau^+\tau^-$)
- NP sensitivity would improve with data \gg LHCb, Belle II, tera-Z (flavor exp. for FCC- hh !?)

Workshop advertisement



Physics at the Flavoured Circular Collider — Next workshop @ CERN: June 16–19
Goal: deepen understanding of the potential of the FCC for flavor physics (2025–2027)

Conclusions

- Flavor physics probes scales $\gg 1$ TeV, sensitivity limited by statistics
New physics could show up any time measurements improve
- There is much to be learned from future measurements at LHCb, Belle II, ATLAS, CMS
- Few tensions with SM — some of these (or others) could soon become decisive
- Interesting challenges both for experiment and theory to maximize sensitivity
Explosion of data always triggered unforeseen developments
- Complementarity between high- p_T and precision probes of new physics will continue
- FCC- ee is a leap from LEP, rich physics program, can be a discovery machine
In flavor physics, the only way to go well beyond Belle II & LHC(b)
- Compelling reasons to aim for the largest data sets that technologies may allow



Extra slides

Theory challenges / opportunities


- **New methods & ideas:** recall that the best α and γ measurements are in modes proposed in light of Belle & BaBar data (i.e., not in the BaBar Physics Book)
 - Better SM upper bounds on $S_{\eta'K_S} - S_{\psi K_S}$, $S_{\phi K_S} - S_{\psi K_S}$, and $S_{\pi^0 K_S} - S_{\psi K_S}$
And similarly in B_s decays, and for $\sin 2\beta_{(s)}$ itself
 - How big can CP violation be in $D^0 - \bar{D}^0$ mixing (and in D decays) in the SM?
 - Many lattice QCD calculations (operators within and beyond SM)
 - Better understanding of inclusive & exclusive semileptonic decays
 - Factorization at subleading order (different approaches), charm loops
 - Can direct CP asymmetries in nonleptonic modes be understood enough to make them “discovery modes”? [$SU(3)$, the heavy quark limit, etc.]
- **We know how to make progress on some + discover new frameworks / methods?**

Another FCNC, $B \rightarrow X_s \gamma$

- Maybe the most complex SM calculation; develop EFTs and RGEs, multi-loop techniques, SCET
Exp. & theor. uncertainty small in different regions
- Extract from global fit short-distance and hadronic parameters (shape functions) fully consistently
[Bernlochner, Lacker, ZL, Stewart, Tackmann, Tackmann, 2007.04320]
- **SIMBA**: Consistent theory across E_γ spectrum
Model-independent treatment of shape fn.

$$|C_7^{\text{incl}} V_{tb} V_{ts}| = (14.77 \pm 0.51_{\text{fit}} \pm 0.59_{\text{theory}} \pm 0.08_{\text{param}}) \times 10^{-3}$$

$$m_b^{1S} = (4.750 \pm 0.027_{\text{fit}} \pm 0.033_{\text{theory}} \pm 0.003_{\text{param}}) \text{ GeV}$$



We find underestimated uncertainty by HFLAV
 \Rightarrow more room for NP

CP violation in D decays and mixing

- CP violation in D decays:

LHCb, Nov. 2011: $\Delta A_{CP} \equiv A_{K^+K^-} - A_{\pi^+\pi^-} = -(8.2 \pm 2.4) \times 10^{-3}$ (I think a stretch in the SM)


LHCb, Mar. 2019: $\Delta A_{CP} = -(1.82 \pm 0.33) \times 10^{-3}$ [1903.08726]

- What is the maximal CPV that could be due to SM?

CKM factors: $|V_{cb}V_{ub}/(V_{cd}V_{ud})| \simeq 7 \times 10^{-4}$



Before measurements, most theory papers stated (assumed) that strong interaction suppresses CPV further

- Can we establish if CP violation in decay or mixing (more “inclusive”) could still probe BSM?



CP violation in $D - \bar{D}$ mixing

- Mixing generated by down quarks
- **SUSY**: up-type squarks in box diagrams, interplay of D & K bounds \Rightarrow alignment, universality, heavy squarks?
- Connections to FCNC top decays
- **Only learned recently**: $x/y = \mathcal{O}(1)$
(Only in 2021 was $\Delta m \neq 0$ established at $> 3\sigma$)
- Very high scales probed, further improvements expected



CP violation in $B_{d,s}$ mixing: $A_{\text{SL}}^{d,s}$


- Only observed in Kaons

$B_{(s)}$: SM suppressed by m_c^2/m_b^2 , may be lifted by BSM

[hep-ph/0202010]

$$A_{\text{SL}} = \frac{\Gamma[\bar{B}^0(t) \rightarrow \ell^+ X] - \Gamma[B^0(t) \rightarrow \ell^- X]}{\Gamma[\bar{B}^0(t) \rightarrow \ell^+ X] + \Gamma[B^0(t) \rightarrow \ell^- X]}$$

Plenty of room between current sensitivity and SM predictions (not yet known if LHCb becomes syst. limited)



- Current status: Exp: $A_{\text{SL}}^d = -(2.1 \pm 1.7) \times 10^{-3}$ $A_{\text{SL}}^s = -(0.6 \pm 2.8) \times 10^{-3}$
SM: $A_{\text{SL}}^d = -(4.7 \pm 0.6) \times 10^{-4}$ $A_{\text{SL}}^s = (2.22 \pm 0.27) \times 10^{-5}$ [1603.07770]
- Unique to Tera-Z: uncertainty $\sim 2.5 \times 10^{-5}$ for both A_{SL}^d and A_{SL}^s , reach SM level

Aside: why not do this in SMEFT?

- Operator analysis (Lorentz invariance, not adding ν_R):

$$(\bar{c}\gamma_\mu P_L b)(\bar{\tau}\gamma^\mu P_L \nu), (\bar{c}\gamma_\mu P_R b)(\bar{\tau}\gamma^\mu P_L \nu), (\bar{c}P_R b)(\bar{\tau}P_L \nu), (\bar{c}P_L b)(\bar{\tau}P_L \nu), (\bar{c}\sigma^{\mu\nu} P_L b)(\bar{\tau}\sigma_{\mu\nu} P_L \nu)$$

Whether b, τ, c in L - or R -handed fields, connection to $b \rightarrow s\tau^+\tau^-$, $b \rightarrow s\nu\bar{\nu}$, $t \rightarrow c\tau^+\tau^-$

2nd and 5th terms can only arise from dim-8 \Rightarrow often neglected

Connection to different generation transitions, only if some flavor structure is imposed

- Semileptonic operators ($l\bar{l}q\bar{q}$) are a large fraction of the operator basis:

SMEFT: 1053 semileptonic operators, 42% of the 2499 parameters of the dim-6 B & L conserving terms in the 3-generation SMEFT (558 CP -even, 495 CP -odd) [\[2402.09535\]](#)

LEET: 1944 semileptonic param's, 54% of the 3631 terms (1017 CP -even, 927 CP -odd)

Similar decay of kaons: $K \rightarrow \pi \nu \bar{\nu}$

- Kaon CPV is at the right level (can fit ϵ_K with KM phase, but ϵ'_K notoriously hard)

- $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ searched for since 1960s (longer than Higgs), sensitive to 100 TeV scale


Irreducible theory uncertainty few % (& $|V_{cb}|^4$)

- **Recently:** first $> 5\sigma$ observation [NA62, 2412.12015]



$$\mathcal{B}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (13.0^{+3.3}_{-3.0}) \times 10^{-11}$$

Consistent with SM ($\approx 8 \times 10^{-11}$), at 1.7σ

- **KOTO:** $\mathcal{B}(K_L \rightarrow \pi^0 \nu \bar{\nu}) < 2 \times 10^{-9}$



Higgs to fermion couplings is what flavor is...



- No constraint yet on origin of 1st generation fermion masses
- FCC- ee can establish role of Higgs in y_c , get close to y_s and y_e