

How Higgs is the Higgs?

- **A study of Vector Boson Scattering at μ TRISTAN**


Risshin Okabe (Kavli IPMU, U Tokyo)

Work in progress

with **Ryuichiro Kitano** (YITP) and **Satoshi Shirai** (IPMU)

Motivation - How Higgs is the Higgs?

An important role of Higgs: **unitarization of WW scatterings**




Deviation of Higgs coupling **enhances** WW scatterings with $O(E^2)$

- ▶ How much does the 125 GeV Higgs unitarize the scatterings?

UV complete amplitudes with an additional scalar

Higgs coupling deviation can be complemented by an additional scalar

$$\mathcal{M}(W_L^+ W_L^+ \rightarrow W_L^+ W_L^+) =$$




Current LHC constraints by indirect Higgs coupling measurements

$$|\delta_h| < 0.084 \quad (95\% \text{ CL})$$

[Lane, Lewis, Sullivan (2024)]


Enhancements of $\sigma(W_L^+ W_L^+ \rightarrow W_L^+ W_L^+)$



μ TRISTAN: $\mu^+ \mu^+$ collider


[Hamada, Kitano, Matsudo, Takaura, Yoshida (2022)]

- **Lepton collider**
 - Clean environment
- **High center-of-mass energy**
 - Probe high-energy growth
- **Cooling techniques of μ^+ beam**
 - High luminosity



Ideal for precise measurement of “Higgsness”

Effects on $\sigma_{\text{tot}}(\mu^+\mu^+ \rightarrow \bar{\nu}\bar{\nu} W^+W^+)$



$W_L W_L$ contribution
 suppressed by
 small parton luminosity

But BSM effect can be $O(1)\%$


Kinematic observables

Four observables

- WW invariant mass M_{WW}
- Scattering angle of WW θ_{WW}
- Decay angles of two W 's $\theta_{1,2}^*$

Final States


- Two hadronic decays
- Semi-leptonic (hadronic decay + lepton decay)



Setup for analysis

- Center-of-mass energy: $\sqrt{s} = 10 \text{ TeV}$
- Integrated luminosity: $L = 10 \text{ ab}^{-1}$
- Cuts
 - ▶ Invariant mass: $M_{WW} > 800 \text{ GeV}$
 - ▶ Pseudo-rapidity: $|\eta| < 1.5$
- Binned χ^2 analysis: 6 bins for M_{WW} , 3 bins for θ_{WW} , 3 bins for $\theta_{1,2}^*$

Results - Sensitivity reach (95% CL)



Summary

- **Higgs is the key to unitarization of Vector Boson Scattering (VBS)**
 - Deviation of hWW coupling can be detected through VBS
- **Same-sign lepton collider $\rightarrow t, u$ -channels**
 - Not affected by resonance peaks
 - Complementary probe to s -channel resonance searches
- **μ TRISTAN is great test site for Higgs measurements!**

Next steps

- **Other unitarization mechanisms**
 - ▶ Heavy vector boson Z' ? Discrimination from scalar case?
- **Estimation of backgrounds**
 - ▶ $ZW, ZZ, q\bar{q}$ channels
- **Effective vector approximation vs Full tree-level calculation**
 - ▶ Particle distribution function may be less precise in high energies
 - ▶ Compare with MadGraph simulation