

# Mid-Internship Progress Report

## KEK–J-PARC Summer Student Program 2026

### Hypernuclear Physics and Detector Calibration

Srushti S. Gaikwad

Manipal Institute of Applied Physics  
Manipal Academy of Higher Education  
India

July 2026

# About Me



## **Srushti S. Gaikwad**

Fourth-Year B.Sc. (Hons.) Physics Student  
Manipal Academy of Higher Education  
Manipal, India

## **Research Interests and Background**

- Worked with HPGe detectors for the pre-final year project
- Strong interest in Gamma ray spectroscopy and Nuclear Physics

# Overview

## Training

- Initial Training at KEK
- Hypernuclear Physics

## Research Activities

- ROOT Simulations
- HPGe Detector Calibration
- Calibration Script Development

## Current Work

- Counter Calibration
- Time-of-Flight Analysis (Ongoing)

# Initial Training at KEK

During the first phase of the internship, I attended lectures given by **Dr. Kyoichiro Ozawa** covering the fundamental concepts required for the project.

Topics included

- Basics of Nuclear Physics
- Basics of Particle Physics
- Radiation Detectors
- Introduction to CERN ROOT

As part of the training, a simulation exercise was performed to generate a uniform decay distribution in momentum vector space using CERN ROOT.

# Uniform Decay Distribution Simulation

- Monte Carlo simulation in CERN ROOT
- Random generation of momentum vectors
- Uniform distribution over solid angle
- Visualization in momentum space

The resulting isotropic momentum distribution demonstrates the uniform generation of momentum vectors in three-dimensional momentum space.




Figure 1: Simulated isotropic momentum distribution generated using CERN ROOT.

# Introduction to Hypernuclear Physics

During the internship, I attended lectures by **Dr. Mifuyu Ukai** introducing the physics motivation behind the experiment.

The topics covered included

- Exotic particles
- Hypernuclei
- Hyperon–nucleon interactions
- The K1.8 beamline experiment at J-PARC

These lectures provided the theoretical background required to understand the ongoing experimental work and subsequent data analysis.

# Exotic Atom

An exotic atom is formed when an electron is replaced by another negatively charged particle.

## Examples

- Muonic atoms
- Pionic atoms
- Kaonic atoms
- Antiprotonic atoms
- $\Xi^-$  atoms

## Applications

- Strong interaction
- Nuclear structure




Figure 2: Illustration of an exotic atom.

# The K1.8 Beamline Experiment

The K1.8 beamline at J-PARC is designed to produce secondary particle beams for hypernuclear physics experiments.


The beamline enables

- Production of secondary  $K^-$  beams
- Momentum analysis using magnetic spectrometers
- Particle identification using dedicated detector systems
- Hypernuclear spectroscopy through the  $(K^-, K^+)$  reaction

In the  $(K^-, K^+)$  reaction, an incoming  $K^-$  beam interacts with a nuclear target to produce a  $\Xi^-$  hyperon, which may subsequently be captured by the nucleus to form a  $\Xi$  hypernucleus. The small production cross section requires high beam intensity together with efficient particle identification and background suppression.



# Experimental Setup of the K1.8 Beamline



**Figure 3:** Schematic layout of the K1.8 beamline showing the production and  $\gamma$ -ray spectroscopy of  $\Xi^-$  hypernuclei.

## Visit to the Hadron Experimental Facility

As part of the internship, I visited the **Hadron Experimental Facility** at J-PARC.

The facility tour, conducted by **Dr. T. Yamamoto**, included an introduction to:

- Beam transport system
- Experimental beamlines
- Detector setup
- Data acquisition system



**Figure 4:** Experimental area of the Hadron Experimental Facility at J-PARC.



## Woods–Saxon Nuclear Density Distribution

One of the first computational tasks was to study the nuclear density distribution of the  $^{12}\text{C}$  nucleus using the Woods–Saxon model.

$$\rho(r) = \frac{\rho_0}{1 + \exp\left(\frac{r-c}{z}\right)}$$

where

- $\rho_0$  : Central nuclear density
- $c$  : Half-density radius
- $z$  : Surface diffuseness parameter

The density distribution was implemented and visualized using CERN ROOT.

# Simulated Nuclear Density

## Simulation

- Carbon-12 nucleus
- Woods–Saxon model
- Numerical implementation in ROOT
- Density calculated as a function of radius




Figure 6: Woods–Saxon nuclear density distribution of the  $^{12}\text{C}$  nucleus.

# Hydrogen-like Exotic Atoms

The radial wavefunction of a hydrogen-like exotic atom is given by

$$\psi(r) \propto r^{n-1} \exp\left(-\frac{r}{na_0}\right)$$

where


- $n$  : Principal quantum number
- $a_0$  : Effective Bohr radius
- $r$  : Distance from the nucleus

The wavefunctions were calculated and visualized for several quantum states using CERN ROOT.

# Radial Wavefunctions

The simulations show that

- Higher quantum numbers correspond to larger orbital radii.
- The probability distribution shifts away from the nucleus.
- Different exotic particles produce different orbital sizes due to their reduced masses.



# Calculated Orbital Radii and Binding Energies

Using the hydrogen-like model, orbital radii and binding energies were calculated for

- Antiprotonic Carbon Atoms
- $\Xi^-$  Carbon Atoms

The calculations demonstrate the influence of reduced mass on atomic structure.

Binding Energy ( $B_e$ ) for  $\Xi^-$  and  $\bar{p}$

| $\Xi^-$ (Xi minus) |            |             |             | $\bar{p}$ (Antiproton) |            |            |             |
|--------------------|------------|-------------|-------------|------------------------|------------|------------|-------------|
| n                  | $r_n$ (fm) | $E_n$ (eV)  | $E_n$ (keV) | n                      | $r_n$ (fm) | $E_n$ (eV) | $E_n$ (keV) |
| 1                  | 3.81       | -1132095.15 | -1132.09515 | 1                      | 5.2        | -829338.48 | -829.33848  |
| 2                  | 15.25      | -283023.79  | -283.02379  | 2                      | 20.8       | -207334.62 | -207.33462  |
| 3                  | 34.31      | -125788.35  | -125.78835  | 3                      | 46.8       | -92148.72  | -92.14872   |
| 4                  | 61.00      | -76755.95   | -76.75595   | 4                      | 83.3       | -51833.66  | -51.83366   |
| 5                  | 95.31      | -45283.81   | -45.28381   | 5                      | 130.1      | -33173.54  | -33.17354   |
| 6                  | 137.24     | -31447.09   | -31.44709   | 6                      | 187.3      | -23037.18  | -23.03718   |

$r_n$ : Radial distance (fm)

$E_n$  (eV): Binding Energy in electron volts



# HPGe Energy Calibration

## Purpose

Establish the energy calibration procedure for a representative HPGe detector using reference  $\gamma$ -ray sources.

## Calibration Procedure

- ① Read detector data from ROOT files.
- ② Construct ADC histograms.
- ③ Fit reference photopeaks with Gaussian functions.
- ④ Determine photopeak centres.
- ⑤ Match ADC channels to known  $\gamma$ -ray energies.




Figure 7: Gaussian fit to a representative  $\gamma$ -ray photopeak for energy calibration.

# Calibrated Energy Spectrum

The photopeak centres obtained from the Gaussian fits were matched with reference  $\gamma$ -ray energies to establish the energy calibration relation for a representative detector.

$$E = A \times ADC + B$$

The above calibration relation was then applied to convert the ADC spectrum into a calibrated energy spectrum.




Figure 8: Calibrated  $\gamma$ -ray energy spectrum of a representative HPGe detector.

# Development of the Calibration Script

## Objective

Automate the energy calibration procedure for multiple detector channels.

## Implementation

- Developed a CERN ROOT script with **Mr. Yuning Hong**.
- Automated the calibration workflow for all 16 detector channels.
- Merged the calibrated spectra from all detector segments into a single spectrum for further analysis.




Figure 9: Calibrated energy spectra obtained for all 16 detector channels.

# Current Progress: Counter Calibration and Time-of-Flight Analysis

## Current Work

- Calibrating scintillation counter signals for the beamline detector system.
- Determining peak centres for each detector segment.
- Extracting calibration parameters for subsequent analysis.
- Preparing calibrated data for Time-of-Flight (TOF) measurements.

## Objective

The calibrated counter data will be used to determine particle time-of-flight, enabling particle identification in the experimental analysis.

*Counter calibration is currently in progress.*

# References

-  W. R. Leo, *Techniques for Nuclear and Particle Physics Experiments*, 2nd Edition, Springer (1994).
-  R. Brun and F. Rademakers, *ROOT – An Object-Oriented Data Analysis Framework*, Nucl. Instrum. Methods Phys. Res. A **389**, 81–86 (1997).
-  Lecture material by Dr. Kyoichiro Ozawa, KEK–J-PARC Summer Student Program (2026).
-  Lecture material by Dr. Mifuyu Ukai, KEK–J-PARC Summer Student Program (2026).



# Physics Information from Antiprotonic X-rays

- The emitted X-rays correspond to transitions between atomic energy levels.
- Strong interaction between the antiproton and the nucleus shifts the lowest atomic energy levels.
- The widths of the X-ray transitions are related to the absorption of the antiproton by the nucleus.
- Precise X-ray spectroscopy provides valuable information on the antiproton–nucleus interaction and serves as a benchmark for theoretical calculations.

# Hyperball-X HPGe Detector Array

- Hyperball-X is a 16-channel high-purity germanium (HPGe) detector array.
- Designed for high-resolution  $\gamma$ -ray and X-ray spectroscopy.
- Used in hypernuclear and exotic atom experiments at the K1.8 beamline of J-PARC.
- High energy resolution enables precise determination of transition energies.




Figure 11: Hyperball-X HPGe detector array used at J-PARC.



## Merged Calibrated Spectrum

- Individual calibrated spectra from all 16 HPGe detectors were merged.
- The combined spectrum provides improved counting statistics for analysis.




Figure 12: Merged calibrated  $\gamma$ -ray energy spectrum obtained by combining all 16 HPGe detector channels