

5d/6d exceptional instantons from trivalent gluing of web diagrams

Hiroataka Hayashi
(林 博貴)

(Tokai University)

Based on the collaboration with

▪ [Kantaro Ohmori \(IAS\)](#)

[\[arXiv:1702.07263\]](#)

Exceptional Groups as Symmetries of Nature on 18th of July at KEK

1. Introduction

- The topological vertex is a powerful tool to compute the all genus topological string amplitudes for toric Calabi-Yau threefolds.

Iqbal 02, Aganagic, Klemm, Marino, Vafa 03
Awata, Kanno 05, Iqbal, Kozcaz Vafa 07

- The full topological string partition function has a physical meaning as the Nekrasov partition function through M-theory on toric Calabi-Yau threefolds.
- We can compute a large class of Nekrasov partition functions regardless of whether the theories have a Lagrangian description or not.

- However there are still many interesting 5d theories to which we had not known how to apply the topological vertex.

Ex.

(1) 5d pure $SO(2N)$ gauge theory

(2) 5d pure E_6, E_7, E_8 gauge theories



ADHM construction is not known

(Nevertheless, some results are known)

Benvenuti, Hanany, Mekareeya 10, Keller, Mekareeya, Song, Tachikawa 11, Gaiotto, Razamat 12, Keller, Song 12, Hananay, Mekareeya, Razamat 12, Cremonesi, Hanany, Mekareeya, Zaffaroni 14, Zafir 15

- In this talk, we will present a powerful prescription of using the topological vertex to compute the partition functions of 5d pure $SO(2N)$, E_6 , E_7 , E_8 gauge theories by utilizing their dual descriptions.
- In fact, the technique can be also applied to 5d theories which arise from a circle compactification of 6d “pure” $SU(3)$, $SO(8)$, E_6 , E_7 , E_8 gauge theories with one tensor multiplet.

1. Introduction
2. 5d gauge theories from string theory
3. A dual description of 5d DE gauge theories
4. Trivalent gluing prescription
5. Applications to 5d theories from 6d
6. Conclusion

2. 5d gauge theories from string theory


- We construct 5d theories with eight supercharges from string theory.
- There are mainly two ways to construct such 5d theories.
 1. M-theory compactification on a non-compact Calabi-Yau three-fold.
 2. (p, q) 5-brane webs.

Witten 96, Morrison Seiberg 96,
Douglas, Katz, Vafa 96

Aharony, Hanany 97,
Aharony, Hanany, Kol 97


1. M-theory compactification on a non-compact Calabi-Yau three-fold.
 - An ADE gauge symmetry is realized by an ADE singularity over a curve.
 - The genus of the curve is related to the number of adjoint hypermultiplets. In this talk we only consider the genus zero case.

- Ex. 5d pure SU(2) gauge theory
→ A_1 singularities over a sphere



← A_1 Dynkin diagram

- Ex. 5d pure $SU(2)$ gauge theory
→ A_1 singularities over a sphere




← A_1 Dynkin diagram



An M2-brane wrapping the fiber sphere gives a W-boson.

- Ex. 5d pure SU(2) gauge theory
→ A_1 singularities over a sphere




← A_1 Dynkin diagram




Non-Abelian SU(2) gauge symmetry is recovered in a limit when the fiber sphere shrinks.

- Ex. 5d pure $SU(2)$ gauge theory
→ A_1 singularities over a sphere




- Ex. 5d pure $SU(2)$ gauge theory
→ A_1 singularities over a sphere



Whole surface shrinks

→ 5d superconformal field theory

- Ex. 5d pure SU(2) gauge theory
→ A_1 singularities over a sphere



Whole surface shrinks

→ 5d superconformal field theory

“UV complete”

2. (p, q) 5-brane webs


- A 5d theory is realized on a worldvolume theory of 5-branes.
- The 5-brane configuration in Type IIB string theory.

	0	1	2	3	4	5	6	7	8	9
D5-brane	×	×	×	×	×	×				
NS5-brane	×	×	×	×	×		×			
(p, q) 5-brane	×	×	×	×	×	angle				

5-brane web


Aharony, Hanany 97,
Aharony, Hanany, Kol 97

Ex. Pure $SU(2)$ gauge theory in a Coulomb branch.




- 6d theory on a 1d space \rightarrow effectively a 5d theory

Ex. Pure $SU(2)$ gauge theory in a Coulomb branch.




- The 5d fixed point is realized when all the particles become massless.



- The two descriptions are in fact dual to each other.
- A 5-brane web in type IIB string theory is dual to a toric diagram of a toric Calabi-Yau threefold in M-theory.


Leung Vafa 97



3. A dual description of 5d DE gauge theories

- We would like to find a dual description of 5d pure $SO(2N)$, E_6 , E_7 , E_8 gauge theories.
- Since we consider D, E gauge groups, we first start from M-theory configurations.
- ADE gauge groups are obtained from ADE singularities over a sphere in a Calabi-Yau threefold

- Ex. 5d pure $SO(2N+4)$ gauge theory
→ D_{N+2} singularities over a sphere




← Dynkin diagram of $SO(10)$

- We can take a different way to see the same geometry for a dual description.

“**fiber-base duality**”


Katz, Mayr, Vafa 97
Aharony, Hanany, Kol 97
Bao, Pomoni, Taki, Yagi 11



- We can take a different way to see the same geometry for a dual description.

“**fiber-base duality**”

Katz, Mayr, Vafa 97
Aharony, Hanany, Kol 97
Bao, Pomoni, Taki, Yagi 11




← base




shrink other spheres




- A schematic picture



- A schematic picture






- A schematic picture




SU(2) gauge theory

- A schematic picture




- The 5d SCFTs may be thought of as “matter” for the $SU(2)$ gauge theory.
- Due to the $SU(2)$ gauge symmetry, each of the 5d SCFTs should have an $SU(2)$ flavor symmetry.
- What are the matter SCFTs?


- Going back to the schematic picture




- Going back to the schematic picture



- Going back to the schematic picture



- Going back to the schematic picture



- In fact, there are two pure $SU(2)$ gauge theories depending on the discrete theta angle θ .

Seiberg 96
Morrison Seiberg 96,
Douglas, Katz, Vafa 96


- The UV completion of the two theories are 5d SCFTs but their flavor symmetries are different.

(i). $\theta = 0 \rightarrow SU(2)$ flavor symmetry (E_1 theory)

(ii). $\theta = \pi \rightarrow U(1)$ flavor symmetry (\tilde{E}_1 theory)


- Therefore, the 5d SCFT should be the E_1 theory.

- It is illustrative to see it from 5-brane webs.
- A 5-brane web for E_1 theory.




pure $SU(2)$

- It is illustrative to see it from 5-brane webs.
- A 5-brane web for E_1 theory.




- It is illustrative to see it from 5-brane webs.
- A 5-brane web for E_1 theory.




pure SU(2)

→
S-dual




- It is illustrative to see it from 5-brane webs.
- A 5-brane web for E_1 theory.




pure SU(2)

→
S-dual




- It is illustrative to see it from 5-brane webs.
- A 5-brane web for E_1 theory.



pure SU(2)


S-dual



perturbative SU(2)


non-Lagrangian

- It is illustrative to see it from 5-brane webs.
- A 5-brane web for E_1 theory.



pure SU(2)


S-dual





- $\widehat{D}_N(SU(2))$ is a 5d SCFT with $(N-1)$ -dimensional Coulomb branch moduli space and has an $SU(2)$ flavor symmetry.


Del Zotto, Vafa, Xie 15

- When the $SU(2)$ flavor symmetry is perturbative the theory is S-dual to a pure $SU(N)$ gauge theory with the CS level N or $-N$.












original picture:






original picture:




→ pure SU(3) gauge theory


- The pure $SU(3)$ gauge theory should have an $SU(2)$ flavor symmetry hence the Chern-Simons level should be 3 or -3 .
- A 5-brane web picture:




- The pure $SU(3)$ gauge theory should have an $SU(2)$ flavor symmetry hence the Chern-Simons level should be 3 or -3 .
- A 5-brane web picture:





→
S-dual




- The pure $SU(3)$ gauge theory should have an $SU(2)$ flavor symmetry hence the Chern-Simons level should be 3 or -3 .
- A 5-brane web picture:





 S-dual



- The geometric picture




- The shrinking limit leads to:




- A duality

pure $SO(10)$ gauge theory



- In general

pure $SO(2N+4)$ gauge theory




- In general


pure $SO(2N+4)$ gauge theory




“trivalent gauging”



- A web-like description




- A web-like description




- We will make use of this picture for the later computations by topological strings.

- In fact, this realization of a duality can be easily extended to pure E_6 , E_7 , E_8 gauge theories.


- In fact, this realization of a duality can be easily extended to pure E_6 , E_7 , E_8 gauge theories.
- Ex. pure E_6 gauge theory




- In fact, this realization of a duality can be easily extended to pure E_6 , E_7 , E_8 gauge theories.
- Ex. pure E_6 gauge theory




- A fiber – base duality




- A fiber – base duality



- A fiber – base duality




- The shrinking limit leads to:




- A duality

Pure E_6 gauge theory




- A web-like picture




- A duality for pure E_7 gauge theory

Pure E_7 gauge theory



- A duality for pure E_8 gauge theory

Pure E_8 gauge theory



4. Trivalent gluing prescription

- We propose a prescription for computing the partition functions of the dual theories which are constructed by the trivalent gauging.
- For that let us consider a simpler case of an $SU(2)$ gauge theory with one flavor.

- The Nekrasov partition function of an SU(2) gauge theory with one flavor is schematically written by

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} Z^{hyper}_{\lambda, \mu}$$

Young diagrams describing the fixed points of U(1) in the U(2) instanton moduli space.

SU(2) vector multiplets

- The Nekrasov partition function of an SU(2) gauge theory with one flavor is schematically written by

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} Z^{hyper}_{\lambda, \mu}$$

Young diagrams describing the fixed points of U(1) in the U(2) instanton moduli space.

SU(2) instanton background

- Therefore, we would like to generalize this expression to

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} Z^{T_1}_{\lambda, \mu} Z^{T_2}_{\lambda, \mu} Z^{T_3}_{\lambda, \mu}$$

Trivalent SU(2) gauging of three 5d SCFTs

- Therefore, we would like to generalize this expression to

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} Z^{T_1}_{\lambda, \mu} Z^{T_2}_{\lambda, \mu} Z^{T_3}_{\lambda, \mu}$$


Trivalent SU(2) gauging of three 5d SCFTs

How can we compute these partition functions?

- However, obtaining the partition functions for the matter theories with an $SU(2)$ instanton background will be difficult from a Lagrangian point of view since the $SU(2)$ flavor symmetry appears non-perturbatively.
- We argue that the topological vertex methods helps us to compute the partition functions of the matter theories with an $SU(2)$ instanton background.

- The topological vertex formalism is kind of a Feynman rule for the topological string amplitude.

Iqbal 02,
 Aganagic, Klemm, Marino, Vafa 03




- For external legs, we assign a trivial Young diagram.

- The full topological string partition function can be computed by summing over Young diagrams of a product of topological vertices.

$$Z_{top} = \sum_{\alpha_i, \beta_i, \gamma_i} \prod_i (-Q_{\alpha_i})^{|\alpha_i|} (-Q_{\beta_i})^{|\beta_i|} (-Q_{\gamma_i})^{|\gamma_i|} C_{\alpha_i \beta_i \gamma_i}$$

- The topological string partition function yields the Nekrasov partition function up to extra factors.

- A naive expectation is that we can simply apply the topological vertex to the web-diagram with non-trivial Young diagrams on the parallel external legs.

$$Z^{\widehat{D}_N(SU(2))}_{\lambda, \mu} \stackrel{?}{=} \text{Diagram}$$


- We propose that the correct prescription is given by dividing it by a half of the $SU(2)$ vector multiplet contribution.

$$Z^{\widehat{D}_N(SU(2))}_{\lambda, \mu} = \left[\text{Diagram 1} \right] / \left[\text{Diagram 2} \right]$$


- Hence when we consider the trivalent $SU(2)$ gauging of three 5d SCFTs, $\widehat{D}_{N_1}(SU(2))$, $\widehat{D}_{N_2}(SU(2))$, $\widehat{D}_{N_3}(SU(2))$, we argue that the partition function is given by

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} \times \underbrace{Z^{\widehat{D}_{N_1}(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_{N_2}(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_{N_3}(SU(2))}_{\lambda, \mu}}_{\text{partition functions of three 5d SCFT matter}}$$

partition functions of three 5d SCFT matter

- With this prescription, it is now straightforward to compute the partition functions of 5d pure $SO(2N+4)$, E_6 , E_7 , E_8 gauge theories.

(1). Pure $SO(2N+4)$ gauge theories




- The partition function:

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} \\ \times Z^{\widehat{D}_N(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu}$$

- We checked that this indeed agrees with the localization result in the unrefined limit until the order Q^8 for the perturbative part and also until the order Q^5 for the one-instanton part and the two-instanton part for the case of $SO(8)$.

(2). Pure E_6 theory



- The partition function

$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} \\ \times Z^{\widehat{D}_3(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_3(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu}$$

- We checked the result in the unrefined limit agrees with the localization result.

Perturbative part : until Q^6

One-instanton part : until Q^2

- The computation for the E_7 and E_8 partition functions is straightforward and we performed non-trivial checks.

Remarks:

1. It is possible to include matter in the vector representation for the $SO(2N+4)$ gauge theory.
2. We can compute the partition function of $SO(2N+3)$ gauge theory by a Higgsing from the partition function of $SO(2N+4)$ gauge theory with vector matter.
3. We can extend the computation to the refined topological vertex. We checked the validity for $SO(8)$.

5. Applications to 5d theories from 6d

- The trivalent gauging method can be also applied to 5d theories which arise from 6d SCFTs on a circle.
- We consider 6d pure $SU(3)$, $SO(8)$, E_6 , E_7 , E_8 gauge theories with one tensor multiplet.
- They are examples of non-Higgsable clusters and important building blocks for constructing general 6d SCFTs.

Morrison, Taylor 12,

Heckman, Morrison Vafa 13

Del Zotto, Heckman, Tomasiello, Vafa 14


Heckman, Morrison Rudelius, Vafa 15

- Those 6d SCFTS can be realized by F-theory compactifications on non-compact elliptically fibered Calabi-Yau threefolds.
- In the case of the pure $SU(3)$, $SO(8)$, E_6 , E_7 , E_8 gauge theories, the geometries have type IV , I_0^* , IV^* , III^* , II^* fibration over a sphere respectively.
- Basically, the fiber spheres form an **affine** Dynkin diagram.


- 5d descriptions for the 6d pure $SO(8)$, E_6 , E_7 , E_8 gauge theories have been already known.

Del Zotto, Vafa, Xie 15


- A 5d description of 6d $SO(8)$ gauge theory without matter:



- Affine Dynkin (6d) vs Dynkin (5d)




6d pure $SO(8)$




5d pure $SO(8)$

- Affine Dynkin (6d) vs Dynkin (5d)



6d SCFT



5d SCFT

- The partition function of the 5d theory is given by


$$Z_{Nek} = \sum_{\lambda, \mu} Q^{|\lambda|+|\mu|} Z^{SU(2)}_{\lambda, \mu} \\ \times Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu} Z^{\widehat{D}_2(SU(2))}_{\lambda, \mu}$$

- The elliptic genus of this 6d SCFT has been computed.


Haghighat, Klemm, Lockhart, Vafa 14

- We checked that the result agrees with the one-string elliptic genus in the unrefined limit until the order $Q^2 Q_4^2$.


- It is straightforward to extend the analysis to the cases of 6d pure E_6 , E_7 , E_8 gauge theories with one tensor multiplet.
- Namely, we extend the Dynkin fibers of E_6 , E_7 , E_8 to the **affine** Dynkin fibers.
- Ex. E_6



- E_7



- E_8



- We computed the partition functions from the trivalent gauging prescription.

- Finally, we consider the 6d pure $SU(3)$ gauge theory with one tensor multiplet.
- The structure of the geometry is different from the previous cases and we start from its geometry.
- The Calabi-Yau threefold for the 6d theory can be realized by an orbifold.

- The orbifold geometry is given by $T^2 \times \mathbb{C}^2 / \Gamma$ with an orbifold action


$$\begin{array}{c} (\omega^2; \omega, \omega) \quad \text{with } \omega^3 = 1 \\ T^2 \quad \mathbb{C} \quad \mathbb{C} \end{array}$$

- The torus becomes a sphere with three fixed points. But there is no singularity over the sphere.
- The fixed point geometry is locally given by $\mathbb{C}^3 / \mathbb{Z}_3$, which is local \mathbb{P}^2 geometry.


- A 5d description can be obtained by considering M-theory on the same Calabi-Yau threefold.

Vafa 96

- Then each of the fixed points gives a 5d SCFT, E_0 theory, coming from the local \mathbb{P}^2 . And three are coupled with each other.




- Local \mathbb{P}^2 geometry




non-Lagrangian

- The Calabi-Yau geometry is given by gluing three local \mathbb{P}^2 geometries.


- Gluing two local \mathbb{P}^2 geometries.




- Gluing two local \mathbb{P}^2 geometries.



- Gluing three local \mathbb{P}^2 geometries



- We can use the same gluing technique to compute the partition function of the SCFT from a local \mathbb{P}^2 geometry.

$$Z^{E_0}_v =$$


The diagram illustrates the gluing technique for computing the partition function. It shows two trivalent vertices connected by a thick black line. The top vertex has three edges extending upwards and to the left. The bottom vertex has three edges extending downwards and to the right. A green arrow labeled v points to the right edge of the top vertex, and another green arrow labeled v points to the right edge of the bottom vertex.

- Then the partition function of the 5d theories from the 6d pure SU(3) gauge theory is given by

$$Z_{Nek} = \sum_{\nu} Q^{|\nu|} Z^{SU(1)}_{\nu} Z^{E_0}_{\nu} Z^{E_0}_{\nu} Z^{E_0}_{\nu}$$




partition function of a resolved conifold

- The elliptic genus of this 6d SCFT has been recently calculated.


Kim, Kim, Park 16

- For comparison we in fact need to perform flop transitions.
- When we take a 5d limit by taking the size of the compactification circle to infinity then the 6d theory reduces to a pure $SU(3)$ gauge theory.

- In the current case, decoupling one local \mathbb{P}^2 reproduces the geometry glued by two local \mathbb{P}^2 .



- In the current case, decoupling one local \mathbb{P}^2 reproduces the geometry glued by two local \mathbb{P}^2 .



- Therefore, we need to perform the flop transition for the partition function obtained from the trivalent $SU(1)$ gauging of three local \mathbb{P}^2 geometries.
- After the flop transition, indeed we found agreement with the elliptic genus of one-string until the order of $Q_1^2 Q_2^2 Q_3^2$.

Remarks:

1. Among the other non-Higgsable clusters, the one with gauge groups $SU(2) \times SO(7) \times SU(2)$ has an orbifold construction. We determined the 5d description and it is again given by the trivalent $SU(2)$ gauging.
2. We can extend the computation to the refined topological vertex. We checked the case of $SO(8)$ until the order $Q Q_1^2 Q_2^2 Q_3^3$ for the one-string part.

6. Conclusion

- We proposed a **new** prescription to compute the partition functions of 5d theories constructed by **trivalent gauging**.
- This method gives the Nekrasov partition functions of **(B)DE** gauge theories in addition to AC.
- Furthermore, we computed the partition functions of 5d theories from circle compactifications of 6d pure $SU(3)$, $SO(8)$, E_6 , E_7 , E_8 gauge theories.