

Anomaly matching in QCD thermal phase transition

Kazuya Yonekura
Tohoku U.

Based on

- [1706.06104] with Hiroyuki Shimizu
- [1901.08188]

Introduction

QCD phase transition is important for cosmology:
Axion abundance etc.

Most radical scenario: [Witten, 1984]

If the phase transition is **first order**, the dark matter might be produced purely by QCD phase transition.
(Several other conditions need to be satisfied.)

The dark matter might be explained by the standard model!

Introduction

Some lattice simulations say that QCD phase transition is **cross-over (i.e. no definite phase transition)**.

But it is not completely settled yet, especially in the limit of small quark masses.

Therefore, it is desirable to study it by methods which do not rely on numerical simulations.

Introduction

A rough version of my claim

(I will explain more precise technical result later.)

If

- Small quark mass approximation is good,
- Large N expansion is good,

then

- QCD phase transition may be naturally first order.

Introduction

Both **small quark mass approximation** and **large N expansion** are qualitatively very good in QCD at zero temperature.

- **Chiral perturbation theory,...**
- **Most mesons as $q\bar{q}$ (rather than $qq\bar{q}\bar{q}$), OZI rule,**
- **Simulation for pure Yang-Mills, ...**

Crossover phase transition may be in tension with those good concepts of QCD and the argument I discuss later.

Contents

1. Introduction

2. 't Hooft Anomaly matching

3. Confinement in finite temperature QCD

4. Results and implications

5. Derivation of Anomaly

6. Summary

't Hooft anomaly

What method do we have to study strong dynamics such as QCD?

't Hooft anomaly matching

UV:

gauge fields + fermions with
global symmetry F



confinement

IR:

???

Anomaly of F in UV = Anomaly of F in IR

't Hooft anomaly in QCD

't Hooft anomaly matching in QCD at zero temperature

In QCD, there exist approximate **chiral symmetry**

$$SU(N_f)_L \times SU(N_f)_R$$

$SU(N_f)_L$: rotate left handed quarks

$SU(N_f)_R$: rotate right handed quarks

Chiral symmetry has the well-known 't Hooft anomaly at zero temperature.

't Hooft anomaly in QCD

UV:

The quarks have the 't Hooft anomaly




confinement

IR:

If there is no chiral fermion,
the chiral symmetry must be spontaneously broken.

't Hooft anomaly in QCD

't Hooft anomaly matching gives an important relation between the two most important concepts in QCD:



How about finite temperature?

The usual anomaly associated to triangle diagram vanishes at finite temperature.

Anomaly at finite temperature

I will argue the existence of a subtler anomaly at finite temperature if we include imaginary chemical potential.

Related works:

[Gaiotto, Kapustin, Komargodski, Seiberg,]
[Tanizaki, Kikuchi, Misumi, Sakai]

→ Misumi-san's talk

Contents

1. Introduction

2. 't Hooft Anomaly matching


3. Confinement in finite temperature QCD

4. Results and implications

5. Derivation of Anomaly

6. Summary

A problem in QCD



We want to study this relation at finite temperature.

However, a well-known problem is that “confinement” is not well-defined in finite temperature QCD because dynamical quarks can screen color fluxes.

Pure Yang-Mills

Let us recall how to define confinement in pure Yang-Mills.

Finite temperature: $Z = \text{tr} e^{-\beta H} \longleftrightarrow R^3 \times S^1$

$\beta = T^{-1}$: inverse temperature

Polyakov loop: $W = \text{tr} P \exp(i \oint_{S^1} A_\mu dx^\mu)$

Wilson loop wrapping on the S^1

Pure Yang-Mills

Intuitively the Polyakov loop behaves as

$$W \sim \exp(-\beta E_q)$$

E_q : energy of a single **probe quark**

Confinement : $E_q \rightarrow \infty$ $W = 0$

Deconfinement : $E_q < \infty$ $W \neq 0$


Order parameter

So the Polyakov loop can be regarded as an order parameter of confinement in **pure-Yang-Mills**.

How about **QCD** with dynamical quarks?

QCD

In QCD, the probe quark energy E_q is always finite.



The Polyakov loop W cannot be used to define confinement phase. Always $W \neq 0$

Imaginary chemical potential

To define confinement rigorously,
I slightly change the problem.

$$\text{tr exp}(-\beta H) \rightarrow \text{tr exp}(-\beta H + i\mu_B B)$$

B : baryon number charge

μ_B : baryon **imaginary chemical potential**

This changes the thermodynamics, but I will argue that the effect of the imaginary chemical potential is subleading in the large N_c expansion.

Imaginary chemical potential

I take

$$\mu_B = \pi$$

[Roberge-Weiss, 1986]

What is special about this value?

All gauge invariant composites have **integer** $B \in \mathbb{Z}$

Mesons: $B = 0$ Baryons: $B = 1$


However, quarks have **fractional** baryon numbers.

Quarks: $B = 1/N_c$

$$\exp(i\pi B) = \begin{cases} \text{real for gauge invariant composites} \\ \text{imaginary for colored quarks} \end{cases}$$


Criterion for confinement

$$W \sim \exp(-\beta E_q + i\pi B)$$



$$\text{Im}(W) \neq 0$$

deconfinement



$$\text{Im}(W) = 0$$

confinement

\mathbb{Z}_2 symmetry for confinement

$$W = \text{tr} P \exp\left(i \oint_{S^1} A_\mu dx^\mu\right)$$

By flipping the direction of integration on S^1 , we get

$$W \rightarrow W^*$$

This is a \mathbb{Z}_2 symmetry.

The order parameter of this \mathbb{Z}_2 is precisely $\text{Im}(W)$

$$\mathbb{Z}_2 : \text{Im}(W) \rightarrow -\text{Im}(W)$$

Definition of confinement

We can summarize the above discussion as follows.

- There exists a \mathbb{Z}_2 symmetry (flipping S^1 direction)
- Imaginary part of the Polyakov loop $\text{Im}(W)$ is charged under the \mathbb{Z}_2
- Confinement and deconfinement are distinguished by

Deconfinement : $\text{Im}(W) \neq 0$ \mathbb{Z}_2 broken

Confinement : $\text{Im}(W) = 0$ \mathbb{Z}_2 unbroken

Remark on imaginary chemical

The effect of imaginary chemical potential is very suppressed in the large N expansion:

$$\frac{\text{effect of } \mu_B}{\text{total free energy}} \sim \frac{N_f}{N_c^3}$$

This follows from the fact that
the baryon charge of quarks is $1/N_c$

Therefore, the situation at $\mu_B = 0$ should be similar to $\mu_B = \pi$ as far as large N expansion is qualitatively good.

Contents

1. Introduction
2. 't Hooft Anomaly matching
3. Confinement in finite temperature QCD
4. Results and implications
5. Derivation of Anomaly
6. Summary

Symmetry and Anomaly

Massless QCD at finite temperature with imaginary chemical potential $\mu_B = \pi$ has (at least) two symmetries:

- Chiral symmetry $SU(N_f)_L \times SU(N_f)_R$
- \mathbb{Z}_2 symmetry

Result : (derivation later)

[KY, 2019]

There exists a mixed 't Hooft anomaly between chiral symmetry and \mathbb{Z}_2 symmetry.

This is a parity anomaly in 3-dimensions.

Symmetry and Anomaly



Result : (derivation later)


[KY, 2019]

There exists a mixed 't Hooft anomaly between chiral symmetry and \mathbb{Z}_2 symmetry.

This is a parity anomaly in 3-dimensions.

Implications to phase transition

Let me discuss the implications of the anomaly to QCD phase transition.



Implications to phase transition

Two critical temperatures:

T_{chiral} : critical temperature for chiral symmetry

$T_{\text{deconfine}}$: critical temperature for \mathbb{Z}_2 symmetry

Let us consider possible scenarios. Either


(1) $T_{\text{deconfine}} > T_{\text{chiral}}$

(2) $T_{\text{deconfine}} < T_{\text{chiral}}$

(3) $T_{\text{deconfine}} = T_{\text{chiral}}$

Scenario 1


Scenario 1: $T_{\text{deconfine}} > T_{\text{chiral}}$



We need complicated massless degrees of freedom to match the anomaly.

Scenario 2


Scenario 2: $T_{\text{deconfine}} < T_{\text{chiral}}$



Chiral symmetry breaking ($q\bar{q}$ condensation) happens in deconfinement phase.

Scenario 3

Scenario 3: $T_{\text{deconfine}} = T_{\text{chiral}}$



It may be natural if the phase transition is **first order** to avoid complicated d.o.f. at the critical temperature,

Natural scenario?

There are many logical possibilities, but **a first order transition at a single critical temperature** may be the most natural scenario.

Otherwise, the 't Hooft anomaly requires either of the following:

- (1) Complicated massless d.o.f. for anomaly matching
- (2) $q\bar{q}$ condensation in deconfinement phase
- (3) Something more complicated

Implication for real QCD

Suppose the phase transition is first order for

$$m_q = 0, \quad \mu_B = \pi$$

Then it is expected to remain first order for

$$m_q \neq 0, \quad \mu_B = 0$$

as far as

$$m_q \ll \Lambda, \quad 1/N_c \ll 1$$

Contents

1. Introduction
2. 't Hooft Anomaly matching
3. Confinement in finite temperature QCD
4. Results and implications
5. Derivation of Anomaly
6. Summary

Reduction from 4 to 3 dim.

Thermodynamics is described by compactification

$$\text{spacetime: } R^4 \rightarrow R^3 \times S^1$$

In the absence of gauge fields,
fermions have **anti-periodic boundary condition**.

$$\Psi(x, \tau + \beta) = -\Psi(x, \tau)$$

τ : coordinate of S^1

β : circumference of S^1

Boundary condition

Gauge fields effectively changes the boundary condition.

$$U = P \exp(i \oint A_\mu dx^\mu)$$

: holonomy of gauge fields around S^1

Effectively

(more precisely in a gauge in which locally $A_4 = 0$)

$$\Psi(x, \tau + \beta) = -U \Psi(x, \tau)$$

Boundary condition

The determinant is

$$\det U = e^{i\mu_B} = -1 \quad (\mu_B = \pi)$$

If U preserves the \mathbb{Z}_2 symmetry of flipping S^1 ,

$$U = \text{diag}(\underbrace{-1, \dots, -1}_K, \underbrace{+1, \dots, +1}_{N_c - K})$$

$$\det U = (-1)^K = -1 \quad : K \text{ is odd.}$$

Boundary condition

$$\Psi(x, \tau + \beta) = -U\Psi(x, \tau)$$

$$U = \text{diag}(-1, \dots, -1, +1, \dots, +1)$$

The diagram shows two horizontal double-headed arrows. The left arrow is red and is positioned below the first set of '-1' terms in the matrix U. Below it is the label 'K'. The right arrow is blue and is positioned below the second set of '+1' terms in the matrix U. Below it is the label 'N_c - K'.

Among N_c color components,


K components: **periodic** condition

$N_c - K$ components: **anti-periodic** condition

This means that $K = \text{odd}$ fermions are massless in 3-dim.

Parity anomaly in 3-dim.

The K -massless fermions in 3d have **parity anomaly**.
This is a mixed anomaly between



Parity in 3d comes from Lorentz symmetry in 4d which flips the S^1 -direction.


This is the \mathbb{Z}_2 -symmetry which I talked about.

Contents

1. Introduction
2. 't Hooft Anomaly matching
3. Confinement in finite temperature QCD
4. Results and implications
5. Derivation of Anomaly
6. Summary

Summary

- There exists a subtle 't Hooft anomaly in finite temperature QCD when an imaginary chemical potential is introduced.



- A first order transition may be the most natural scenario of QCD phase transition if large N expansion and small quark mass approximation are qualitatively good.

