

Big Bounce Baryogenesis

Neil D. Barrie

Kavli IPMU (WPI), UTIAS, University of Tokyo

February 18, 2020

arxiv:2001.04773

Matter-antimatter Asymmetry

The asymmetry is described quantitatively by,

$$\eta = \frac{n_b - n_{\bar{b}}}{s} \simeq 8.5 \times 10^{-11}$$

The Sakharov Conditions

- 1 Baryon number violation
- 2 \mathcal{C} and \mathcal{CP} violation
- 3 Period of non-equilibrium

Standard Model $\rightarrow \eta_{sm} \sim 10^{-18}$.

Inflationary dilution \Rightarrow Typically generated during or after reheating.

Inflationary Baryogenesis

- Pseudoscalar inflaton coupled to $F\tilde{F}$,
- Generation of Chern-Simons number from rolling of scalar field,

$$\frac{\phi}{\Lambda} Y_{\mu\nu}^a \tilde{Y}^{a\mu\nu}, \quad \frac{\phi}{\Lambda} W_{\mu\nu}^a \tilde{W}^{a\mu\nu}$$


- Can seed galactic magnetic fields, and generate gravitational wave signatures.
- Y suffers from uncertainties of EWPT and MHD.

Ekpyrotic Bounce

Ekpyrotic Contraction: $a = (\epsilon H_b t)^{\frac{1}{\epsilon}} = (\epsilon H_b |\tau|)^{\frac{1}{\epsilon-1}}$ with $H = -\frac{1}{\epsilon|\tau|}$

Require $\epsilon \geq 3$, leading to very slow contraction for large ϵ .

$$\rho = \frac{\rho_k}{a^2} + \frac{\rho_m}{a^3} + \frac{\rho_r}{a^4} + \frac{\rho_a}{a^6} + \dots + \frac{\rho_\phi}{a^{2\omega}} + \dots$$



Single Field Ekpyrotic Bounce

Achieved if the φ is fast-rolling down a negative exponential potential,

$$V(\varphi) \approx -V_0 e^{-\sqrt{2\epsilon} \frac{\varphi}{M_p}} \quad \text{and} \quad \epsilon = \frac{3}{2}(1 + \omega) .$$

Scaling solution,

$$\varphi \simeq M_p \sqrt{\frac{2}{\epsilon}} \ln(-\sqrt{\epsilon V_0 T} / M_p) \quad \text{and} \quad \varphi' \simeq \sqrt{\frac{2}{\epsilon}} \frac{M_p}{T} .$$

- Solves the problem of the rapid growth of anisotropies.
- Anisotropic instabilities which may arise are suppressed.
- Permits trajectories which are attractors.
- Predict small r .
- Can produce large non-gaussianities.

The Model and Gauge Field Dynamics

Lagrangian terms in contracting background:

$$\mathcal{L} = -\frac{1}{4}\eta^{\mu\rho}\eta^{\nu\sigma}F_{\mu\nu}F_{\rho\sigma} - \frac{\varphi}{8\Lambda}\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu}F_{\rho\sigma} ,$$

Satisfying the Sakharov Conditions

- 1 Anomalous currents,
- 2 Pseudoscalar coupling to Chern-Simons terms,
- 3 Ekpyrotic Contracting phase.

The potential for the Ekpyrotic Scalar ,

$$V(\varphi) \approx -V_0 e^{-\sqrt{2\epsilon}\frac{|\varphi|}{M_p}} .$$

Field Quantisation and Mode Functions

- Derive equations of motion F_i , in weak field limit,
- Solving for circularly polarised wave modes ($\alpha = +, -$),

$$F_i = \int \frac{d^3\vec{k}}{(2\pi)^{3/2}} \sum_{\alpha} \left[G_{\alpha}(\tau, k) \epsilon_{i\alpha} \hat{a}_{\alpha} e^{i\vec{k}\cdot\vec{x}} + G_{\alpha}^*(\tau, k) \epsilon_{i\alpha}^* \hat{a}_{\alpha}^{\dagger} e^{-i\vec{k}\cdot\vec{x}} \right] .$$

- Thus,

$$G_{\pm}'' + \left(k^2 \mp \frac{2\kappa k}{-\tau} \right) G_{\pm} = 0 , \quad \text{where } \kappa = \frac{M_p}{\sqrt{2\epsilon\Lambda}}$$

with solutions,

$$G_{\pm} = \frac{e^{-ik\tau}}{\sqrt{2k}} e^{\pm\pi\kappa/2} U(\pm i\kappa, 0, 2ik\tau)$$

- Interested in the exponentially enhanced positive frequency modes.

Scenario 1: Baryon Number from $\varphi W\tilde{W}$

Related to the baryon number,

$$\partial_\mu (\sqrt{-g} j_B^\mu) = \frac{3g_2^2}{32\pi^2} \epsilon^{\mu\nu\rho\sigma} W_{\mu\nu}^a W_{\rho\sigma}^a = \frac{3g_2^2}{16\pi^2} \partial_\mu (\sqrt{-g} K^\mu) .$$

Baryon number density at τ ,

$$\begin{aligned} \frac{n_B(\tau)}{a(\tau)^3} &\simeq \frac{9g_2^2}{8\pi^4} \int_\mu^{2\epsilon\kappa|H|} k^3 (|G_+(\tau)|^2 - |G_-(\tau)|^2) dk \\ &\simeq \frac{9g_2^2}{16\pi^4} (-\epsilon|H|)^3 C(\kappa) , \end{aligned}$$

where

$$C(\kappa) \sim 0.007 \frac{e^{2\pi\kappa}}{\kappa^4}, \text{ for } \kappa > 1 .$$

Can now calculate the asymmetry parameter.

Generated Baryon Asymmetry

- No significant entropy production after reheating ($s \simeq \frac{2\pi^2}{45} g^* T_{\text{rh}}^3$),
- Heavy majorana ν_R and reheating temperature $T_{\text{rh}} > 10^{12}$ GeV
- Evaluating $n_B(\tau)$ near the bounce,

$$\frac{\eta_B}{\eta_B^{\text{obs}}} \simeq 2 \cdot 10^5 C(\kappa) \left(\frac{\epsilon |H_c|}{T_{\text{rh}}} \right)^3 .$$


Require $\rho_{CS}^W(\tau) \ll 3M_p^2 H^2$, to ensure gauge fields do not effect the background evolution,

$$M_p \gg \sqrt{\frac{6C(\kappa)}{\pi}} \epsilon^2 |H_c| .$$

For $H_c = H_b$

Successful Baryogenesis requires,

$$|H_b| \simeq \frac{2 \cdot 10^{14} \text{ GeV}}{\epsilon^2 C(\kappa)^{2/3}} \quad \text{and} \quad T_{\text{rh}} \simeq \frac{10^{16} \text{ GeV}}{\epsilon C(\kappa)^{1/3}}$$




Energy density constraint,

$$1 \ggg 10^{-4} \kappa^{2/3} e^{-\pi\kappa/3} \quad \text{or} \quad 1 \ggg \epsilon \left(\frac{T_{\text{rh}}}{10^{25} \text{ GeV}} \right)$$

For $|H_c| < |H_b|$

Successful Baryogenesis requires,

$$|H_c| = 2 \cdot 10^{-2} \frac{T_{\text{rh}}}{\epsilon C(\kappa)^{1/3}},$$



A κ -independent bound on H_c for the energy density constraint,

$$\frac{H_c}{H_b} > \epsilon \frac{T_{\text{rh}}}{4 \cdot 10^{22} \text{ GeV}}$$

Scenario 1: Summary

- Ample parameter space for which successful Baryogenesis,
- Performed simplified calculation using the linearised approximation,
- This approximation will likely break down for large κ . A more detailed analysis is required.
- Indicates that Baryogenesis may be possible through this mechanism.

Scenario 2: Hypermagnetic Field Helicity from $\varphi Y \tilde{Y}$

The hypermagnetic field generated is,

$$B_{\text{rh}}(\tau_b)^2 = \frac{1}{2\pi^2} \int_{\mu}^{2\epsilon\kappa|H_c|} k^4 (|G_+(\tau)|^2 - |G_-(\tau)|^2) dk$$

which can be expressed as,

$$B_{\text{rh}}(\tau_b) \simeq \frac{1}{2\pi} (\epsilon H_c)^2 \sqrt{\frac{2C(\kappa)}{\kappa}}.$$

with the approximate correlation length of these magnetic fields,

$$\lambda_{\text{rh}}(\tau_b) \simeq \frac{4\pi\kappa}{\epsilon|H_c|}.$$

These follow known evolution from T_{rh} to the EWPT.

Case 2: $\varphi Y \tilde{Y}$

The baryon asymmetry parameter produced at the EWPT is within the range,

$$C(\kappa) \left(\frac{H_c}{H_b}\right)^3 \left(\frac{\epsilon^2 |H_b|}{10^{17} \text{ GeV}}\right)^{3/2} < \frac{\eta_B}{\eta_B^{obs}} < C(\kappa) \left(\frac{H_c}{H_b}\right)^3 \left(\frac{\epsilon^2 |H_b|}{1.5 \cdot 10^{15} \text{ GeV}}\right)^{3/2}$$

Requiring $\rho_{CS}^Y(\tau) \ll 3M_p^2 H^2$,

$$M_p \gg \sqrt{\frac{2C(\kappa)}{\pi}} \epsilon^2 |H_c|.$$

Taking the parameters required for successful Baryogenesis gives,

$$2.4 \cdot 10^{-17} \text{ G} < B_p^0 < 2 \cdot 10^{-16} \text{ G} \quad \text{and} \quad 7 \cdot 10^{-4} \text{ pc} < \lambda_p^0 < 6 \cdot 10^{-3} \text{ pc}$$


Below constraints, but unable to explain Blazar observations.

$$H_c = H_b$$

$$\frac{1.5 \cdot 10^{15} \text{ GeV}}{C(\kappa)^{2/3}} < \epsilon^2 |H_b| < \frac{10^{17} \text{ GeV}}{C(\kappa)^{2/3}} .$$

hence considering the maximum value, Eq. (13) becomes,

$$C(\kappa) \gg 10^{-9} ,$$





Upper bound corresponds to lower bound uncertainty on asymmetry generation, and vice versa. $\kappa = 1, 3, 5$

$$|H_c| < |H_b|$$

$$\frac{H_c}{H_b} \gg \frac{\epsilon T_{\text{rh}}}{2.5 \cdot 10^{20} \text{ GeV}}, \quad \text{and} \quad \frac{H_c}{H_b} \gg \frac{\epsilon T_{\text{rh}}}{10^{23} \text{ GeV}}.$$

Parameter space for successful Baryogenesis, considering a reheating temperature of $T_{\text{rh}} \sim 10^{15} \text{ GeV}$.



For the lower bound and upper bound on asymmetry generation, respectively, for $\kappa = 1, 3, 5$.

Conclusion and Future Work

- Ekpyrotic phase, induced by a fast-rolling pseudoscalar field,
- Chern-Simons number density generated through coupling to φ
- Successfully produce η_B^{obs} through both the Y_i and W_i gauge fields.
- For Y_i , also source galactic magnetic fields, similar to inflation case.

Future Investigations

- Detail the bounce dynamics and back-reaction effects,
- Possibly gravitational wave signature.