

Modeling Evolution of Dark Matter Substructure and Annihilation Boost

Reference: Phys.Rev.D97., 123002

Nagisa Hiroshima (Univ. of Tokyo, KEK)

Collaboration with

S.Ando (Univ. of Amsterdam), T.Ishiyama (Chiba Univ.)

Contents

- **Motivations**
- **Modeling**
- **Applications**
- **Summary**


Motivations

WIMP Dark Matter


- Naturally explains the relic abundance with weak scale cross section

$$\langle\sigma v\rangle \sim 3 \times 10^{-26} \text{cm}^3 \text{s}^{-1}$$

- WIMP DM mass should be $m_{\text{DM}} \sim \mathcal{O}(1)\text{GeV} - \mathcal{O}(1)\text{TeV}$
- predict formation of small scale structures down to $10^{-12} - 10^{-3} M_{\odot}$




DM search with γ -rays



$$\phi_\gamma = \frac{1}{2} \frac{1}{4\pi} \frac{\langle \sigma v \rangle}{m_{\text{DM}}^2} \int \frac{dN_\gamma}{dE_\gamma} dE_\gamma \cdot$$

$$\int_{l.o.s} \rho_{\text{DM}}^2 ds$$

subhalo boost



$$\phi_\gamma = \frac{1}{2} \frac{1}{4\pi} \frac{\langle \sigma v \rangle}{m_{\text{DM}}^2} \int \frac{dN_\gamma}{dE_\gamma} dE_\gamma \cdot (1+B) \int_{l.o.s} \rho_{\text{DM}}^2 ds$$

subhalos on l.o.s boost the annihilation signals


Estimates of subhalo boost

1312.1729

- ◆ $M_{\text{halo}} \sim 10^{-6} - 10^{16} M_{\odot}$

- ◆ $z \sim 0 - 10$


**Previous works adopted
extrapolation of results from
numerical simulations**





Needs for physical, wide-coverage modelings

Modeling


The situation





1, collapse



2, accretion



3, Tidal Stripping



4, subhalo on our l.o.s

Evolution of subhalos


1. Formation

2. Accretion

3. Tidal Stripping


Assumptions:

- NFW profile with truncation for host & subhalo
- mass-loss occurs in the first orbit of each subhalo



$$\dot{m} = \frac{m - m(r_t)}{T_r}$$

Tidal Stripping



a single power-law in 20 orders of magnitude


calculations of boost factors

- ✦ **host evolutions**
- ✦ **mass accretion history**
- ✦ **NFW parameters after tidal stripping**

Boost factor $B = \frac{\sum_i w_i \rho_{s,i}^2 r_{s,i}^3}{\rho_{s,\text{host}}^2 r_{s,\text{host}}^3}$


Applications

Mass fraction of subhalo



Good agreements with N-body result


Boost Factor



Annihilation signal is boosted by factors

Update on IGRB limit

DM + DM \rightarrow b \bar{b}




Improving IGRB constraint

Summary

Summary

- **We modeled tidal stripping of subhalos in an analytical way covering more than 20 orders of mass ranges**
- **Subhalos in our model show good agreements with those in N-body simulations**
- **As applications, we calculate boost factor of DM annihilation signals**
- **DM annihilation signals can be boosted up to a factor of 10 in cluster scales**

Boost Factor



Annihilation signal is boosted by factors

Tidal stripping

- **potential of the host**

$$\Phi(R) = -\frac{GM_{\text{host}}}{R_{\text{vir}}} \frac{\ln \left[1 + c_{\text{vir}}^{\text{host}} R/R_{\text{vir}} \right]}{f(c_{\text{vir}})^{\text{host}} R/R_{\text{vir}}}$$

- **orbital period of subhalo**

$$T_r = 2 \int_{R_p}^{R_a} \frac{dR}{\sqrt{2 \left[E - \Phi(R) \right] - L^2/R^2}}$$

- **tidal mass of subhalo**

$$(r_t/R_p)^3 = \frac{m(r_t)}{M_{<R_p}} \left(2 + \left(L^2/R_p GM_{<R_p} \right) - \left(d \ln M(R)/d \ln R \right) |_{R_p} \right)^{-1}$$

- **mass-loss rate**

$$\dot{m} = [m - m(r_t)]/T_r$$

